

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number  
**WO 01/20003 A2**

(51) International Patent Classification<sup>7</sup>: **C12N 15/54**,  
15/11, 9/12, C12Q 1/68, A01K 67/027, G01N 33/68

(21) International Application Number: **PCT/EP00/09896**

(22) International Filing Date:  
11 September 2000 (11.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
99402236.6 10 September 1999 (10.09.1999) EP  
00401388.4 18 May 2000 (18.05.2000) EP

(71) Applicant (for all designated States except US):  
**INSTITUT NATIONAL DE LA RECHERCHE  
AGRONOMIQUE (INRA)** [FR/TR]; 147, rue de l'Uni-  
versité, F-75007 Paris (FR).

(71) Applicants and

(72) Inventors: **ANDERSSON, Leif** [SE/SE]; Bergagatan 30,  
S-752 39 Uppsala (SE). **LOOFT, Christian** [DE/DE];  
Mittelweg 3 c, 24802 Bokelholm (DE). **KALM, Ernst**  
[DE/DE]; Schmalholt 1, 24239 Achterwehr (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MILAN, Denis**  
[FR/FR]; 3 bis, chemin du Tricou, F-31670 Labège (FR).  
**ROBIC, Annie** [FR/TR]; 33, rue des Capitouls, F-31650

Saint-Orens-De-Gameville (FR). **ROGEL-GAILLARD, Claire** [FR/FR]; 156, rue Léon Maurice Nordinant,  
F-75013 Paris (FR). **IANNUCCELLI, Nathalie** [FR/FR];  
7, boulevard des Alouettes, F-31320 Castanet-Tolosan  
(FR). **GELLIN, Joël** [FR/FR]; 8, allée des Amazones,  
F-31320 Auzeville (FR). **LE ROY, Pascale** [FR/FR]; 32,  
avenue Saint Marc, F-91300 Massy (FR). **CHARDON, Patrick** [FR/FR]; 17, rue de Petite Fontaine, F-91430  
Vauhallan (FR).

(74) Agents: **VIALLE-PRESLES, Marie-José** et al.; Cabinet  
Ores, 6, avenue de Messine, F-75008 Paris (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE). OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

Without international search report and to be republished  
upon receipt of that report.

[Continued on next page]

(54) Title: VARIANTS OF THE GAMMA CHAIN OF AMPK, DNA SEQUENCES ENCODING THE SAME, AND USES THEREOF



(57) Abstract: The invention concerns variants of the gamma chain of vertebrate AMP-activated kinase (AMPK), as well as nucleic acid sequences encoding said variants and use thereof for the diagnosis or treatment of dysfunction of energy metabolisms.



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

VARIANTS OF THE GAMMA CHAIN OF AMPK, DNA SEQUENCES  
ENCODING THE SAME, AND USES THEREOF.

The present invention relates to new variants  
of the  $\gamma$  chain of AMP-activated protein kinase (AMPK), to  
5 genes encoding said variants and to uses thereof.

AMPK has a key role in regulating the energy  
metabolism in the eukaryotic cell (HARDIE et al., Annu.  
Rev. Biochem., 67, 821-855, 1998; KEMP et al., TIBS, 24,  
22-25, 1999). Mammalian AMPK is a heterotrimeric complex  
10 comprising a catalytic  $\alpha$  subunit and two non-catalytic  $\beta$   
and  $\gamma$  subunits that regulate the activity of the  $\alpha$   
subunit. The yeast homologue (denoted SNF1) of this  
enzyme complex is well characterised; it comprises a  
catalytic chain (Snf1) corresponding to the mammalian  $\alpha$   
15 subunit, and regulatory subunits: Sip1, Sip2 and Gal83  
correspond to the mammalian  $\beta$  subunit, and Snf4  
correspond to the mammalian  $\gamma$  subunit. Sequence data show  
that AMPK homologues exist also in *Caenorhabditis elegans*  
and *Drosophila*.

20 It has been observed that mutations in yeast  
SNF1 and SNF4 cause defects in the transcription of  
glucose-repressed genes, sporulation, thermotolerance,  
peroxisome biogenesis, and glycogen storage.

In the mammalian cells, AMPK has been proposed  
25 to act as a "fuel gauge". It is activated by an increase  
in the AMP:ATP ratio, resulting from cellular stresses  
such as heat shock and depletion of glucose and ATP.  
Activated AMPK turns on ATP-producing pathways (e.g.  
fatty acid oxidisation) and inhibits ATP-consuming  
30 pathways (e.g. fatty acid and cholesterol synthesis),  
through phosphorylation of the enzymes acetyl-CoA  
carboxylase and hydroxymethylglutaryl-CoA (HMG-CoA)  
reductase. It has also been reported to inactivate in  
vitro glycogen synthase, the key regulatory enzyme of  
35 glycogen synthesis, by phosphorylation (HARDIE et al.,

1998, *supra*); however, whether glycogen synthase is a physiological target of AMPK *in vivo* remained unclear.

Several isoforms of the three different AMPK subunits are present in mammals. In humans, *PRKAA1* on human chromosome (HSA) 5p12 and *PRKAA2* on HSA1p31 respectively encode isoforms  $\alpha 1$  and  $\alpha 2$  of the  $\alpha$  subunit, *PRKAB1* on HSA12q24.1 and *PRKAB2* (not yet mapped) respectively encode isoforms  $\beta 1$  and  $\beta 2$  of the  $\beta$  subunit, and *PRKAG1* on HSA12q13.1 and *PRKAG2* on HSA7q35-q36 respectively encode isoforms  $\gamma 1$  and  $\gamma 2$  of the  $\gamma$  subunit (OMIM database, <http://www.ncbi.nlm.nih.gov/omim/>, July 1999). HARDIE et al., [1998, *supra*] also mention the existence of a third isoform ( $\gamma 3$ ) of the  $\gamma$  subunit of AMPK but do not provide any information about it. Analysis of the sequences of these  $\gamma$  subunits shows that they are essentially composed of four cystathione  $\beta$  synthase (CBS) domains whose function is unknown. No phenotypic effect resulting from a mutation in either of the AMPK subunits has yet been documented.

On the other hand, it has been observed that most Hampshire pigs have a high intramuscular glycogen concentration. In these pigs, glycogenolysis which occurs after slaughtering leads to an important decrease of the pH, resulting in acid meat having a reduced water-holding capacity and giving a reduced yield of cured cooked ham.

The locus (named *RN*) associated with high muscular content of glycogen was first identified by family segregation analysis of phenotypic data from Hampshire pigs (LE ROY et al., Genet. Res., 55, 33-40, 1990). A fully dominant allele, *RN*, correlated with high glycogen content occurs at a high frequency in most Hampshire populations while pigs from other breeds are assumed to be homozygous for the normal, recessive *rn*<sup>+</sup> allele. Subsequent studies showed that *RN* carriers have a large increase (about 70%) of glycogen in skeletal muscle

but not in liver (MONIN et al., in 38<sup>th</sup> ICoMST, Clermont-Ferrand, FRANCE, 1992).

The large difference in glycogen content between RN and rn<sup>+</sup> pigs leads to marked differences in meat quality and technological yield (ENFÄLT et al., J. Anim. Sci., 75, 2924-2935, 1997). The RN allele is therefore of considerable economical significance in the pig industry and most breeding companies would like to reduce or eliminate this dominant mutation.

The RN phenotype can be determined by measuring the glycolytic potential in muscle biopsies from live animals, or after slaughter (MONIN et al., Meat Science, 13, 49-63, 1985). However, this method has severe limitations for application in practical breeding programs. The accuracy of the test is not 100%: as there is some overlap in the phenotypic distribution of RN and rn<sup>+</sup>, the test is not able to distinguish RN/RN homozygotes and RN/rn<sup>+</sup> heterozygotes. Further, the sampling of muscle biopsies on live animals is invasive and costly.

Thus, there is a strong need for the development of a simple diagnostic DNA test for the RN locus. Moreover, the dramatic phenotypic effect of the RN gene in pigs implies that this gene has an important role in the regulation of carbohydrate metabolism in skeletal muscle in other vertebrates, in particular mammals.

Skeletal muscle and liver are the two major reservoirs of glycogen in mammals and the observation of an increased muscular glycogen while liver glycogen is normal suggests that the RN<sup>-</sup> phenotype maybe due to a mutation in a gene expressed in muscle but not in liver. The inventors have previously reported that the RN gene is located on pig chromosome 15 (MILAN et al., Mamm. Genome, 7, 47-51, 1996; MARIANI et al., Mamm. Genome, 7, 52-54, 1996; LOOFT et al., Genetics Selection Evolution, 28, 437-442, 1996). They have now discovered that the RN

allele is associated with a non-conservative mutation in a gene encoding a new muscle-specific isoform of the AMP-activated protein kinase (AMPK)  $\gamma$  chain.

The various aspects of the present invention  
5 are based upon the discovery and characterisation of this mutation and the identification and isolation of the mutant gene.

According to the invention it is shown that a mutation in a  $\gamma$  chain of AMPK results in an altered  
10 regulation of carbohydrate metabolism, demonstrating that AMPK is an essential component of said metabolism. It is also provided a nucleic acid sequence encoding a muscle-specific isoform of the  $\gamma$  chain of AMPK. Thus it is provided means to regulate carbohydrate metabolism, more  
15 specifically to detect and/or correct potential or actual dysfunctions of the regulation of carbohydrate metabolism, in particular in skeletal muscle.

The invention provides a polypeptide comprising an amino acid sequence having at least 70%  
20 identity or at least 85% similarity, preferably 80% identity or at least 90% similarity, more preferably at least 90% identity or at least 95% similarity, and still more preferably at least 95% identity or at least 99% similarity, with the polypeptide SEQ ID NO: 2. The  
25 invention also provides an isolated nucleic acid sequence encoding said polypeptide, as well as the complement of said nucleic acid sequence.

Said polypeptide represents a new muscle-specific isoform of the  $\gamma$  chain of AMPK, and will also be  
30 hereinafter referred as Prkag3; the gene encoding said polypeptide will also be hereinafter referred as PRKAG3.

According to a preferred embodiment of the invention, said polypeptide comprises an amino acid sequence having at least 75% identity, preferably at  
35 least 80% identity with the polypeptide SEQ ID NO: 28.

"Identity" of a sequence with a reference sequence refers to the percent of residues that are the same when the two sequences are aligned for maximum correspondence between residues positions. A polypeptide having an amino acid sequence having at least X% identity with a reference sequence is defined herein as a polypeptide whose sequence may include up to 100-X amino acid alterations per each 100 amino acids of the reference amino acid sequence. Amino acids alterations include deletion, substitution or insertion of consecutive or scattered amino acid residues in the reference sequence.

"Similarity" of a sequence with a reference sequence refers to the percent of residues that are the same or only differ by conservative amino acid substitutions when the two sequences are aligned for maximum correspondence between residues positions. A conservative amino acid substitution is defined as the substitution of an amino acid residue for another amino acid residue with similar chemical properties (e.g. size, charge or polarity), which generally does not change the functional properties of the protein. A polypeptide having an amino acid sequence having at least X% similarity with a reference sequence is defined herein as a polypeptide whose sequence may include up to (100-X) non-conservative amino acid alterations per each 100 amino acids of the reference amino acid sequence. Non-conservative amino acids alterations include deletion, insertion, or non-conservative substitution of consecutive or scattered amino acid residues in the reference sequence.

For instance:

\* searching the "GenBank nr" database using BLASTp (ALTSCHUL et al., Nucleic Acids Res., 25, 3389-3402, 1997) with default settings and the whole sequence

SEQ ID NO: 2 as a query, the higher percents of identity or similarity with SEQ ID NO: 2 were found for:

- $\gamma$ 1 subunit of human AMPK: 65% identity or 82% similarity (score: 399);
- 5       -  $\gamma$ 1 subunit of rat AMPK: 65% identity or 82% similarity (score: 399);
- $\gamma$ 1 subunit of murine AMPK: 64% identity or 80% similarity (score: 390);
- $\gamma$  subunit of Drosophila AMPK: 53% identity or 75% similarity (score: 332);
- 10       - Yeast Snf4: 33% identity or 56% similarity (score: 173);

\* searching the "GenBank nr" database using BLASTp with default settings and the whole sequence  
15 SEQ ID NO: 28 as a query, the higher percents of identity or similarity were found for:

- $\gamma$ 1 subunit of human AMPK: 64% identity or 80% similarity (score: 403);
- $\gamma$ 2 subunit of human AMPK: 62% identity or 83% similarity (score: 425);
- 20       -  $\gamma$ 1 subunit of rat AMPK: 61% identity or 77% similarity (score: 404);
- $\gamma$ 1 subunit of murine AMPK: 63% identity or 79% similarity (score: 394);
- 25       -  $\gamma$  subunit of Drosophila AMPK: 52% identity or 76% similarity (score: 340).

Polypeptides of the invention include for instance any polypeptide (whether natural, synthetic, semi-synthetic, or recombinant) from any vertebrate  
30 species, more specifically from birds, such as poultry, or mammals, including bovine, ovine, porcine, murine, equine, and human, and comprising, or consisting of, the amino acid sequence of either:

- a functional Prkag3; or
- 35       - a functionally altered mutant of Prkag3.



"Functional" refers to a protein having a normal biological activity. Such a protein may comprise silent mutations inducing no substantial change in its activity, and having no noticeable phenotypic effects.

5 Non-limitative examples of functional Prkag3 are:

- a porcine Prkag3 comprising at least the sequence represented in the enclosed sequence listing under SEQ ID NO: 2; this includes, for instance the polypeptide SEQ  
10 ID NO: 28;
- a human Prkag3 comprising at least the sequence represented in the enclosed sequence listing under SEQ ID NO: 4; this includes for instance the polypeptide SEQ  
15 ID NO: 30.

The invention also includes splice variants of Prkag3: for instance, the nucleotide sequence SEQ ID NO: 27, and the corresponding amino-acid sequence SEQ ID NO: 28 on one hand, and the nucleotide sequence SEQ ID  
20 NO: 31 and the corresponding amino-acid sequence SEQ ID NO: 32 on the other hand represent two different splice variants of porcine Prkag3.

A "functionally altered mutant" of a protein comprises one or several mutations inducing a change in  
25 its activity. Such mutations include in particular deletions, insertions, or substitutions of amino acid residues in a domain essential for the biological activity of said protein. They may result for instance in a partial or total loss of activity, or conversely in an  
30 increase of activity, or in an impairment of the response to regulatory effectors. Deletions, insertions, or non-conservative substitutions are more likely to result in a critical effect on the biological activity; however conservative substitutions may also induce a noticeable  
35 effect, if they occur at an important position of an active site of the protein.

Non-limitative examples of functionally altered mutants of Prkag3 are:

- the R41Q variant resulting from the non-conservative substitution of an arginine residue in position 41 of SEQ ID NO: 2 or SEQ ID NO: 4 by a glutamine residue (this substitution results in an important increase of the glycogen content, inducing an increased glycolytic potential of the skeletal muscle);
- the V40I variant resulting from the substitution of a valine residue in position 40 of SEQ ID NO: 2 or SEQ ID NO: 4 by an isoleucine residue (this substitution results in a decrease of the glycogen content and thus of the glycolytic potential of the skeletal muscle).

These substitutions occur inside a portion of the first CBS domain that is highly conserved between Prkag3 and the previously known isoforms of the  $\gamma$  subunit of AMPK.

Residue numbers for Prkag3 refer to the amino acid numbering of SEQ ID NO: 2 or SEQ ID NO: 4. Alignment of human and porcine Prkag3 sequences with previously known  $\gamma 1$  and  $\gamma 2$  isoforms is shown in Figure 3.

The invention also provides mutants of Prkag3 which may for instance be obtained by deletion of part of a Prkag3 polypeptide. Said mutants are generally functionally altered. They may have an identity with the overall Prkag3 sequence lower than 70%. However, the identity of the non-deleted sequences of said mutants, when aligned with the corresponding Prkag3 sequences and more specifically with the corresponding sequences from SEQ ID NO: 2, should remain higher than 70%. Said mutants may for instance result from the expression of nucleic acid sequences obtained by deletion or insertion of a nucleic acid segment, or by a punctual mutation introducing a nonsense codon, in a nucleic acid sequence encoding a functional Prkag3.

The invention also provides a functionally altered mutant of a  $\gamma$  subunit of AMPK, wherein said mutant comprises at least one mutation responsible for said functional alteration located within the first CBS domain, and preferably within the region thereof aligned with the region spanning from residue 30 to residue 50 of SEQ ID NO:2 or SEQ ID NO:4. Said mutation may result from the insertion, deletion, and/or substitution of one amino-acid or of several amino-acids, adjacent or not.

More preferably the mutation is located within the region aligned with the region spanning from residue 35 to residue 45 of SEQ ID NO:2 or SEQ ID NO:4, for instance within the region spanning from residue 65 to residue 75 of the  $\gamma 1$  isoform.

According to a particular embodiment, said mutation is a non-conservative substitution, preferably a R $\rightarrow$ Q substitution. According to another particular embodiment, said mutation is a conservative substitution, preferably a V $\rightarrow$ I substitution.

Advantageously, the mutation is located at a residue corresponding to residue 41 of SEQ ID NO:2 or SEQ ID NO:4, for instance in the case of the  $\gamma 1$  isoform, at residue 70, or at a residue corresponding to residue 40 of SEQ ID NO:2 or SEQ ID NO:4, for instance in the case of the  $\gamma 1$  isoform, at residue 69.

The invention also provides a heterotrimeric AMPK wherein the  $\gamma$  subunit consists of a polypeptide of the invention.

The invention also provides isolated nucleic acid sequences encoding any of the above-defined functional or functionally altered Prkag3 or functionally altered mutants of a  $\gamma$  subunit of AMPK, and nucleic acid sequences complementary of any one of these nucleic acid sequences.

This includes particularly any isolated nucleic acid having the sequence of any of the naturally

occurring alleles of a *PRKAG3* gene, as well as any isolated nucleic acid having the sequence of an artificial mutant of a *PRKAG3* gene, provided that said nucleic acid does not consist of the EST GENBANK  
5 AA178898.

This also includes any isolated nucleic acid having the sequence of a natural or artificial mutant of a *PRKAG1* or a *PRKAG2* gene, wherein said mutant encodes a functionally altered  $\gamma 1$  or  $\gamma 2$  subunit of AMPK as defined  
10 above.

Nucleic acids of the invention may be obtained by the well-known methods of recombinant DNA technology and/or of chemical DNA synthesis. These methods also allow to introduce the desired mutations in a naturally  
15 occurring DNA sequence.

Examples of nucleic acids encoding naturally occurring alleles of a *PRKAG3* gene are represented by SEQ ID NO: 1, which encodes a naturally occurring allele of the porcine gene and SEQ ID NO: 3, which encodes a  
20 naturally occurring allele of the human gene. These sequences may be used to generate probes allowing the isolation of *PRKAG3* from other species or of other allelic forms of *PRKAG3* from a same species, by screening a library of genomic DNA or of cDNA.

The invention also includes genomic DNA  
25 sequences from any vertebrate species, more specifically from birds, such as poultry, or mammals, including in particular bovine, ovine, porcine, murine, equine, and human, comprising at least a portion of a nucleic acid  
30 sequence encoding a polypeptide of the invention, preferably a portion of a *PRKAG3* gene, and up to 500 kb, preferably up to 100 kb of a 3' and/or of a 5' adjacent genomic sequence.

Such genomic DNA sequences may be obtained by  
35 methods known in the art, for instance by extension of a nucleic acid sequence encoding a polypeptide of the

invention, employing a method such as restriction-site PCR (SARKAR et al., PCR Methods Applic., 2, 318-322, 1993), inverse PCR (TRIGLIA et al., Nucleic Acids Res., 16, 8186, 1988) using divergent primers based on a Prkag3 coding region, capture PCR (LAGERSTROM et al., PCR Methods Applic., 1, 111-119, 1991), or the like.

The invention also includes specific fragments of a nucleic acid sequence encoding a polypeptide of the invention, or of a genomic DNA sequence of the invention as well as nucleic acid fragments specifically hybridising therewith. Preferably these fragments are at least 15bp long, more preferably at least 20bp long.

"Specific fragments" refers to nucleic acid fragments having a sequence that is found only in the nucleic acids sequences encoding a polypeptide of the invention, and is not found in nucleic acids sequences encoding related polypeptides of the prior art. This excludes the nucleic acid fragments that consist of a sequence shared with one of the known PRKAG1 or PRKAG2 genes.

"Specifically hybridising fragments" refers to nucleic acid fragments which can hybridise, under stringent conditions, only with nucleic acid sequences encoding a polypeptide of the invention, without hybridising with nucleic acid sequences encoding related polypeptides of the prior art. This excludes the nucleic acid fragments that consist of the complement of a sequence shared with one of the known PRKAG1 or PRKAG2 genes.

Nucleic acid fragments that consist of the EST GENBANK AA178898 or the EST GENBANK W94830 or the complements thereof are also excluded.

Said specific or specifically hybridising nucleic acid fragments may for example be used as primers or probes for detecting and/or amplifying a nucleic acid sequence encoding a polypeptide of the invention. The

invention encompasses set of primers comprising at least one primer consisting of a specific or specifically hybridising nucleic acid fragment as defined above.

The invention also provides recombinant  
5 vectors comprising a nucleic acid sequence encoding a polypeptide of the invention. Vectors of the invention are preferably expression vectors, wherein a sequence encoding a polypeptide of the invention is placed under control of appropriate transcriptional and translational  
10 control elements. These vectors may be obtained and introduced in a host cell by the well-known recombinant DNA and genetic engineering techniques.

The invention also comprises a prokaryotic or eukaryotic host cell transformed by a vector of the  
15 invention, preferably an expression vector.

A polypeptide of the invention may be obtained by culturing the host cell containing an expression vector comprising a nucleic acid sequence encoding said polypeptide, under conditions suitable for the expression  
20 of the polypeptide, and recovering the polypeptide from the host cell culture.

A heterotrimeric AMPK wherein the  $\gamma$  subunit consists of a polypeptide of the invention may be obtained by expressing, together or separately, a nucleic  
25 acid sequence encoding a polypeptide of the invention, a nucleic acid sequence encoding an  $\alpha$  subunit, and a nucleic acid sequence encoding a  $\beta$  subunit, and reconstituting the heterotrimer.

The polypeptides thus obtained, or immunogenic  
30 fragments thereof may be used to prepare antibodies, employing methods well known in the art. Antibodies directed against the whole Prkag3 polypeptide and able to recognise any variant thereof may thus be obtained. Antibodies directed against a specific epitope of a  
35 particular variant (functional or not) of Prkag3 or antibodies directed against a specific epitope of a

functionally altered mutant having a mutation in the first CBS domain of a  $\gamma$  subunit of AMPK, and able to recognise said variant or functionally altered mutant may also be obtained.

5           As shown herein, mutations in a  $\gamma$  subunit of AMPK, and particularly mutations in the first CBS domain of a  $\gamma$  subunit of AMPK are likely to cause disorders in the energy metabolism (e.g. diabetes, obesity) in vertebrates, including humans. Further, mutations in the  
10 first CBS domain or other parts of the *PRKAG3* gene are likely to cause disorders in the muscular metabolism leading to diseases such as myopathy, diabetes and cardiovascular diseases.

          The present invention provides means for  
15 detecting and correcting said disorders.

          More specifically, the present invention is directed to methods that utilise the nucleic acid sequences and/or polypeptidic sequences of the invention for the diagnostic evaluation, genetic testing and  
20 prognosis of a metabolic disorder.

          For example, the invention provides methods for diagnosing of metabolic disorders, more specifically carbohydrate metabolism disorders, and preferably disorders correlated with an altered, in particular an  
25 excessive, glycogen accumulation in the cells, resulting from a mutation in a gene encoding a  $\gamma$  subunit of AMPK, wherein said methods comprise detecting and/or measuring the expression of a functionally altered *PRKAG3* gene, or of a functionally altered mutant of a  $\gamma$  subunit of AMPK  
30 having a mutation within the first CBS domain in a nucleic acid sample obtained from a vertebrate, or detecting a mutation in the *PRKAG3* gene or in a sequence encoding the first CBS domain of a  $\gamma$  subunit of AMPK in the genome of a vertebrate suspected of having such a  
35 disorder.

According to a preferred embodiment of the invention, the disorder is correlated with an altered, in particular an excessive, glycogen accumulation in the muscular cells and results from the expression of a  
5 functionally altered *PRKAG3* gene.

The expression of a functionally altered *Prkag3*, or of a functionally altered mutant of a  $\gamma$  subunit of AMPK having a mutation within the first CBS domain may be detected or measured using either polyclonal or  
10 monoclonal antibodies specific for the functionally altered polypeptides of the invention, as defined above. Appropriate methods are known in the art. They include for instance enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell  
15 sorting (FACS).

The nucleotide sequences of the invention may be used for detecting mutations in the *PRKAG3* gene or in a sequence encoding the first CBS domain of a  $\gamma$  subunit of AMPK, by detection of differences in gene sequences or in  
20 adjacent sequences between normal, carrier, or affected individuals.

The invention provides a process for detecting a mutation in the *PRKAG3* gene or in a sequence encoding the first CBS domain of a  $\gamma$  subunit of AMPK wherein said  
25 process comprises:

- obtaining a nucleic acid sample from a vertebrate;
- checking the presence in said nucleic acid sample of a nucleic acid sequence encoding a mutant *Prkag3*, or a mutant of a  $\gamma$  subunit of AMPK having a mutation within  
30 the first CBS domain, as defined above.

According to a preferred embodiment of the invention there is provided a method for detecting a nucleic acid sequence comprising a mutation in the *PRKAG3* gene or in a sequence encoding the first CBS domain of a  $\gamma$   
35 subunit of AMPK wherein said process comprises:

- obtaining a nucleic acid sample from a vertebrate;



- contacting said nucleic acid sample with a nucleic acid probe obtained from a nucleic acid of the invention and spanning said mutation, under conditions of specific hybridisation between said probe and the mutant sequence to be detected;
- detecting the hybridisation complex.

Preferably, the process of the invention further comprises, prior to hybridisation, PCR amplification from the nucleic acid sample, of a sequence comprising at least the portion of the *PRKAG3* sequence or of the sequence encoding the first CBS domain of the  $\gamma$  subunit of AMPK wherein the mutation is to be detected.

Methods allowing the specific hybridisation of a probe only with a perfectly matching complementary sequence, and useful for the detection of punctual mutations are known in the art. They include for instance Allele Specific PCR (GIBBS, Nucleic Acid Res., 17, 2427-2448, 1989), Allele Specific Oligonucleotide Screening (SAIKI et al., Nature, 324, 163-166, 1986), and the like.

A mutation in the *PRKAG3* gene may also be detected through detection of polymorphic markers closely linked to said mutation.

The invention also provides means for identifying said polymorphic markers, and more specifically polymorphic markers comprised within a genomic DNA sequence comprising at least a portion of a *PRKAG3* gene, and up to 500 kb, preferably 300 kb, more preferably up to 100 kb of a 3' and/or of a 5' adjacent sequence.

Said polymorphic markers may be obtained for instance, by screening a genomic DNA library from a vertebrate with a probe specific for the *PRKAG3* gene, in order to select clones comprising said nucleic acid sequence and flanking chromosomal sequences, and identifying a polymorphic marker in said flanking chromosomal sequences. The allele(s) of a polymorphic

marker associated with a given mutant allele of the *PRKAG3* gene may also easily be identified by use of a genomic DNA library from an individual wherein the presence of said mutant allele has previously been  
5 detected by hybridisation with a nucleic acid probe of the invention.

Polymorphic markers include for instance, single nucleotide polymorphisms (SNP), microsatellites, insertion/deletion polymorphism and restriction fragment  
10 length polymorphism (RFLP). These polymorphic markers may be identified by comparison of sequences flanking the *PRKAG3* gene obtained from several individuals. Microsatellites may also be identified by hybridisation with a nucleic acid probe specific of known  
15 microsatellite motifs.

Once a polymorphic marker has been identified, a DNA segment spanning the polymorphic locus may be sequenced and a set of primers allowing amplification of said DNA segment may be designed.

20 The invention also encompasses said DNA primers.

Detection of a mutation in the *PRKAG3* gene may be performed by obtaining a sample of genomic DNA from a vertebrate, amplifying a segment of said DNA spanning a  
25 polymorphic marker by polymerase chain reaction using a set of primers of the invention, and detecting in said amplified DNA the presence of an allele of said polymorphic marker associated with said mutation.

By way of example, polymorphic markers which  
30 may be obtained according to the invention, and DNA primers allowing the detection of polymorphic markers closely linked to the *RN* allele of porcine *PRKAG3* gene are listed in Table 1 hereinafter.

According to a preferred embodiment of the  
35 invention, the vertebrate is a mammal, preferably a farm animal and more preferably a porcine, and the mutation to

be detected produces a functionally altered Prkag3. The detection of said mutation allows to predict whether said mammal or the progeny thereof is likely to have an intramuscular glycogen concentration higher or lower than the average. An example of such a mutation produces a functionally altered Prkag3 having a R41Q substitution, and resulting in an increased glycogen content in the skeletal muscle.

Another example of such a mutation produces a functionally altered Prkag3 having a V40I substitution, and resulting in a decreased glycogen content in the skeletal muscle. In farm animals having such a mutation, glycogenolysis which occurs after slaughtering is less important than in normal animals, resulting in a higher pH and in a potential better quality of the meat.

The present invention also includes kits for the practice of the methods of the invention. The kits comprise any container which contains at least one specific fragment of a nucleic acid sequence of the invention, or at least one nucleic acid fragment able to specifically hybridise with a nucleic acid sequence of the invention. Said nucleic acid fragment may be labelled. The kits may also comprise a set of primers of the invention. They may be used in conjunction with commercially available amplification kits. They may also include positive or negative control reactions or markers, molecular weight size markers for gel electrophoresis, and the like.

Other kits of the invention may include antibodies of the invention, optionally labelled, as well as the appropriate reagents for detecting an antigen-antibody reaction. They may also include positive or negative control reactions or markers.

The invention further provides means for modulating the expression of vertebrate genes encoding a  $\gamma$  subunit of AMPK, and more specifically of the PRKAG3 gene

and/or the synthesis or activity of the products of said genes.

A purified AMPK heterotrimer comprising wild-type or mutant Prkag3 subunit, or a functionally altered mutant  $\gamma$  subunit having a mutation in the first CBS domain, may be used for screening *in vitro* compounds able to modulate AMPK activity, or to restore altered AMPK activity. This may be done, for instance, by:

- measuring the binding of the compound to said heterotrimer, using for example high-throughput screening methods; or,

- measuring changes in AMPK kinase activity, using for example high-throughput screening methods.

High throughput screening methods are disclosed, for instance, in "High throughput screening: The Discovery of Bioactive Substances", J.P. DEVLIN (Ed), MARCEL DEKKER Inc., New York (1997).

Nucleic acids of the invention may be used for therapeutic purposes. For instance, complementary molecules or fragments thereof (antisense oligonucleotides) may be used to modulate AMPK activity, more specifically in muscular tissue.

Also, a nucleic acid sequence encoding a functional Prkag3 may be used for restoring a normal AMPK function.

Transformed cells or animal tissues expressing a wild-type or mutant Prkag3, or a functionally altered mutant of a  $\gamma$  subunit of AMPK as defined above, or expressing an AMPK comprising said mutant Prkag3, or said functionally altered mutant of a  $\gamma$  subunit of AMPK, may be used as *in vitro* model for elucidating the mechanism of AMPK activity or for screening compounds able to modulate the expression of AMPK.

The screening may be performed by adding the compound to be tested to the culture medium of said cells or said tissues, and measuring alterations in energy

metabolism in said cells or said tissues using methods such as measurements of glucose concentrations (levels), glucose uptake, or changes of the ATP/AMP ratio, glycogen or lipid/protein content.

5           The invention provides animals transformed with a nucleic acid sequence of the invention.

In one embodiment, said animals are transgenic animals having at least a transgene comprising a nucleic acid of the invention.

10           In another embodiment, said animals are knockout animals. "Knockout animals" refers to animals whose native or endogenous *PRKAG3* alleles have been inactivated and which produce no functional *Prkag3* of their own.

15           In light of the disclosure of the invention of DNA sequences encoding a wild-type or mutant *Prkag3*, or a functionally altered mutant of a  $\gamma$  subunit of AMPK, transgenic animals as well as knockout animals may be produced in accordance with techniques known in the art,  
20   for instance by means of *in vivo* homologous recombination.

Suitable methods for the preparation of transgenic or knock-out animals are for instance disclosed in: *Manipulating the Mouse Embryo*, 2<sup>nd</sup> Ed., by  
25   HOGAN et al., Cold Spring Harbor Laboratory Press, 1994; *Transgenic Animal Technology*, edited by C. PINKERT, Academic Press Inc., 1994; *Gene Targeting: A Practical Approach*, edited by A.L. JOYNER, Oxford University Press, 1995; *Strategies in Transgenic Animal Science*, edited y  
30   G.M. MONASTERSKY and J.M. ROBL, ASM Press, 1995; *Mouse Genetics: Concepts and Applications*, by Lee M. SILVER, Oxford University Press, 1995.

These animals may be used as models for metabolic diseases and disorders, more specifically for  
35   diseases and disorders of glycogen metabolism in muscle. For instance they may be used for screening test

molecules. Transgenic animals may thus be used for screening compounds able to modulate AMPK activity. Knockout animals of the invention may be used, in particular, for screening compounds able to modulate energy metabolism, more specifically carbohydrate metabolism, in the absence of functional Prkag3.

The screening may be performed by administering the compound to be tested to the animal, and measuring alterations in energy metabolism in said animal using methods such as glucose tolerance tests, measurements of insulin levels in blood, changes of the ATP/AMP ratio, glycogen or lipid/protein content in tissues and cells.

Transgenic or knock-out farm animals with modified meat characteristics or modified energy metabolism may also be obtained.

The present invention will be further illustrated by the additional description which follows, which refers to examples of obtention and use of nucleic acids of the invention. It should be understood however that these examples are given only by way of illustration of the invention and do not constitute in any way a limitation thereof.

#### **EXAMPLE 1: ISOLATING THE PRKAG3 GENE**

We have screened a porcine Bacterial Artificial Chromosome (BAC) library (ROGEL-GAILLARD et al., Cytogenet and Cell Genet, 851, 273-278, 1999) and constructed a contig of overlapping BAC clones across the region of pig chromosome 15 harbouring the RN gene. These BAC clones were in turn used to develop new genetic markers in the form of single nucleotide polymorphisms (SNPs) or microsatellites (MS) as described in Table 1 below.

Table 1

	Name of marker	BAC clone	Primer sequences	Size of PCR product (bp)	Marker type <sup>a</sup>	Alleles <sup>b</sup>
1	H3	115B9, 156E6, 361B4, 90A9	F: 5'-GGAATTTCAAGTCAGCCAAC-3' (SEQ ID NO: 5) R: 5'-CTTCAAAAGACCGTGCTACT-3' (SEQ ID NO: 6)	114 - 138	MS	114, 126, 128, 132*, 134*, 136, 138
2	MS982H1	982H11	F: 5'-CTGGGAACCTCTATATGCTG-3' (SEQ ID NO: 7) R: 5'-TAGGGAATACAAATCACAG-3' (SEQ ID NO: 8)	114 - 157	MS	114, 140, 142*, 144, 146, 150, 158
3	MS479L3	479L3, 297D7, 852B5, 153B5	F: 5'-CTCCAGCTCACAGGATGACA-3' (SEQ ID NO: 9) R: 5'-GTTCTGCAGCTTAGCATCTATCC-3' (SEQ ID NO: 10)	150 - 164	MS	150*, 160, 162, 164
4	MS987M3	997F12	F: 5'-GAAGTATCCTGGCTTCTGA-3' (SEQ ID NO: 11) R: 5'-GTTCTCCAGGTTCCAGACATCCAC-3' (SEQ ID NO: 12)	138 - 160	MS	138, 144, 152, 154, 160*
5	MS482H8	482E7	F: 5'-GCTTCTGCTGCCCTACTT-3' (SEQ ID NO: 13) R: 5'-GTTCTAAGTTCTACTGTAGACACC-3' (SEQ ID NO: 14)	78 - 90	MS	78, 80, 88*, 90
6	MS337H2	808G10, 947E5, 337G11	F: 5'-CCAAGCTGTGGTGGCTGAAT-3' (SEQ ID NO: 15) R: 5'-CAGCACAGCAGTGCCACCTA-3' (SEQ ID NO: 16)	145 - 165	MS	145, 149, 155, 161*, 165*
7	MS127B1	127G6, 134C9	F: 5'-CAAACCTCTTAGGCGGTG-3' (SEQ ID NO: 17) R: 5'-GTTCTGGAACCTCCATAGCCATGG-3' (SEQ ID NO: 18)	94 - 108	MS	94, 100, 108*, 114
8	CMKAR2	128A3, 337G11, 808G10, 947E7, 1110H12	F: 5'-AGGGTGGATGGTAGGCTTCA-3' (SEQ ID NO: 19) R: 5'-GTCTCGCTCCTGAAGGAAGT-3' (SEQ ID NO: 20)	208	SNP	112A*, 112T*, 158A*, 158G 176A*, 176G
9	127G63	127G6, 134C9, 170D7, 1030A5, 1088F2	F: 5'-AGTCACGTGGCCATGCTATC-3' (SEQ ID NO: 21) R: 5'-CTCAACTGGATTGAGTCAGT-3' (SEQ ID NO: 22)	409	SNP	234A*, 234C
10	VIL1		F: 5'-TTGGCGCACTGTATTCT-3' (SEQ ID NO: 23) R: 5'-AGGCAAGGAAGAGCACAG-3' (SEQ ID NO: 24)	270	SNP	90T, 90G, 120A, 120G, 166C, 166T
11	NRAMP1	315F7, 530A6, 651C12, 1088F2, 1095H3	F: 5'-AGCCGTGGCATCGTTGG-3' (SEQ ID NO: 25) R: 5'-AGAAGGAGACAGACAGGGCGA-3' (SEQ ID NO: 26)	1300	RFLP (Syl)	1: 100+1200 bp 2: 100+200+1000 bp

\*MS=microsatellite; SNP=single nucleotide polymorphism.

<sup>b</sup>Microsatellite alleles are designated according to the length of the amplified fragment while SNPs are denoted according to the polymorphic nucleotide. Alleles associated with the *RN* allele are marked with an asterisk.

The new markers were used together with some previously described markers to construct a high-resolution linkage map. Standard linkage analysis using pedigree data comprising about 1,000 informative meioses for segregation at the RN locus made it possible to exclude RN from the region proximal to MS479L3 and distal to microsatellite Sw936. Linkage Disequilibrium (LD) analysis was done with the same markers and a random sample of 68 breeding boars from the Swedish Hampshire population, scored for the RN phenotype by measuring glycogen content in muscle. The results of LD analysis using the DISMULT program (TERWILLIGER, Am. J. Hum. Genet., 56, 777-787, 1995) are shown in Figure 1. They reveal a sharp LD peak around the markers MS127B1 and SNP127G63. These markers appeared to show complete linkage disequilibrium with the RN allele, i.e. RN was associated with a single allele at these two loci. The most simple interpretation of this finding is that the RN mutation arose on a chromosome carrying these alleles and that the two markers are so closely linked to the RN locus that the recombination frequency is close to 0%. The two markers are both present on the overlapping BAC clones 127G6 and 134C9 suggesting that the RN gene may reside on the same clone or one of the neighbouring clones.

A shot-gun library of the BAC clone 127G6 was constructed and more than 1,000 sequence reads were collected giving about 500,000 base pair random DNA sequence from the clone. The data were analysed and sequence contigs constructed with the PHRED, PHRAP and CONSED software package (University of Washington Genome Center, <http://bozeman.mbt.washington.edu>). The sequence data were masked for repeats using the REPEATMASKER software (<http://ftp.genome.washington.edu/cgi-bin/RepeatMasker>) and BLAST searches were carried out using the NCBI web site (<http://www.ncbi.nlm.nih.gov>).



Three convincing matches to coding sequences were obtained. Two of these were against human cDNA sequences/genes, KIAA0173 described as being similar to pig tubulin-tyrosine ligase and located on HSA2q (UniGene cluster Hs.169910, <http://www.ncbi.nlm.nih.gov/UniGene/>) and CYP27A1 located on HSA2q33-ter (UniGene cluster Hs.82568). The results strongly suggested that the pig coding sequences are orthologous to these human genes as it is well established that the RN region is homologous to HSA2q33-36 (ROBIC et al., Mamm. Genome, 10, 565-568, 1999). However, none of these sequences appeared as plausible candidate genes for RN. The third coding sequence identified in BAC 127G6 showed highly significant sequence similarity to various AMP-activated protein kinase  $\gamma$  sequences including the yeast SNF4 sequence. The cDNA sequence of this gene was determined by RT-PCR and RACE analysis using muscle mRNA from an  $rn^+/rn^+$  homozygote. This sequence is shown in Figure 2 and in the enclosed sequence listing under SEQ ID NO: 1.

Legend of Figure 2:  
5' UTR: 5' untranslated region  
3' UTR: 3' untranslated region  
CDS: coding sequence  
\*\*\*: stop codon  
'-': identity to master sequence  
'.': alignment gap

The frame of translation was determined on the basis of homology to other members in the protein family and assuming that the first methionine codon in frame is the start codon. The polypeptidic sequence deduced on this basis is shown in the enclosed sequence listing under SEQ ID NO: 2.

The complete nucleotidic sequence of pig PRKAG3 cDNA is shown in the enclosed sequence listing under SEQ ID NO: 27 and the complete polypeptidic

sequence is shown in the enclosed sequence listing under SEQ ID NO: 28 and in Figure 3.

Figure 3 shows an amino acid alignment constructed with the CLUSTAL W program (THOMPSON et al., Nucleic Acids Research, 22, 4673-4680, 1994) with representative AMPK  $\gamma$  sequences in the nucleotide databases.

#### Legend of Figure 3:

##### Sequences used:

- 10 HumG1: Genbank U42412
  - MusG1: Genbank AF036535
  - HumG2: Human PRKAG2 (Genbank AJ249976)
  - PigG3: pig PRKAG3 (this study)
  - HumG3: human PRKAG3 (this study)
  - 15 Dros: *Drosophila* (Genbank AF094764)
  - SNF4 (yeast): Genbank M30470
- Both the PRKAG2 and *Drosophila* sequences have longer aminoterminal regions but they do not show significant homology to the aminoterminal region of PRKAG3 and were
- 20 not included.

##### Abbreviations:

- \*: stop codon
  - '-': identity to master sequence
  - ..': alignment gap
- 25 The four CBS domains are overlined and the position of the RN<sup>o</sup> mutation is indicated by an arrow.

Table 2 below shows the amino acid (above diagonal) and nucleotide sequence (below diagonal) identities (in %) among mammalian, *Drosophila* and yeast

30 AMPKG/SNF4 sequences. In the case of pig PRKAG3 and human PRKAG3, the identities were calculated referring to the portions thereof represented respectively by SEQ ID NO: 1 and SEQ ID NO: 3, for the nucleotide sequences, and by SEQ ID NO: 2 and SEQ ID NO: 4, for the amino acid

35 sequences.

TABLE 2

	PigG3	HumG3	HumG1	RatG1	MusG1	HumG2	Dros	SNF4
PigG3	-	97.0	64.2	64.2	63.9	62.6	53.2	34.0
HumG3	90.7	-	63.6	63.6	63.6	62.6	53.5	34.4
HumG1	64.2	64.5	-	96.7	96.3	75.6	60.9	33.5
RatG1	65.8	65.8	88.0	-	97.4	75.3	61.1	33.5
MusG1	65.3	64.8	87.2	92.8	-	74.6	61.7	33.5
HumG2	61.6	61.6	68.1	67.8	65.9	-	63.1	34.5
Dros	58.4	58.4	59.0	59.3	59.0	60.0	-	36.2
SNF4	44.0	44.2	45.4	44.6	45.3	45.7	44.8	-

Figure 4 shows a Neighbor-Joining phylogenetic tree constructed with the PAUP software (SWOFFORD, *Phylogenetic analysis using parsimony (and other methods)*, Sinauer Associates, Inc. Publishers, Sunderland, Massachusetts, 1998) using yeast SNF4 as outgroup; support for branch orders obtained in bootstrap analysis with 1,000 replicates are indicated, scales of tree is indicated at the bottom. The result showed that the pig gene located in the RN region is distinct from mammalian PRKAG1 and PRKAG2 isoforms and most likely orthologous to a human gene represented by the human EST sequence AA178898 (GenBank) derived from a muscle cDNA library. This gene is herein denoted PRKAG3 since it is the third isoform of a mammalian AMP-activated protein kinase  $\gamma$  characterised so far.

The cDNA sequence of this gene was determined by RT-PCR and 5'RACE analysis using human skeletal muscle cDNA (Clontech, Palo Alto, CA). This sequence is shown in Figure 2 and in the sequence listing under SEQ ID NO: 3. The deduced polypeptidic sequence having 97% identity with the porcine sequence SEQ ID NO: 2 (cf. Table 2) is shown on Figure 2 and in the sequence listing under SEQ ID NO: 4.

The complete cDNA sequence is also shown in the enclosed sequence listing under SEQ ID NO: 29; the deduced polypeptidic sequence is shown in the enclosed sequence listing under SEQ ID NO: 30 and in Figure 3.

Using the high resolution human TNG radiation hybrid panel : (<http://shgc-www.stanford.edu/RH/TNGindex.html>) we mapped the human homologs of *PRKAG3*, *CYP27A1* and *KIAA0173*, all present in the porcine BAC127G6. The three  
5 genes are also very closely linked in the human genome. *PRKAG3* was mapped at a distance of 33 cR<sub>50.000</sub> from *KIAA0173* and 52 cR<sub>50.000</sub> from *CYP27A1*, with lod score support of 6.8 and 4.5, respectively.

The established role of AMPK in regulating  
10 energy metabolism, including glycogen storage, and its location in the region showing maximum linkage disequilibrium made *PRKAG3* a very strong candidate gene for RN. This was further strengthened by hybridisation analysis of a human multiple tissue northern blots  
15 (CLONTECH, Palo Alto, CA) using human *PRKAG1* (IMAGE clone 0362755 corresponding to GenBank entry AA018675), human *PRKAG2* (IMAGE clone 0322735 corresponding to GenBank entry W15439) and a porcine *PRKAG3* probe. The results are shown in Figure 5.

20 Legend of Figure 5:

H: Heart, B: Brain, Pl: Placenta, L: Lung,  
Li: Liver, M: Skeletal muscle, K: Kidney, Pa: Pancreas,  
S: Spleen, Th: Thymus, P: Prostate, T: Testis, O: Ovary,  
I: Small intestine, C: Colon (mucosal lining),  
25 PBL: Peripheral Blood Leukocyte.

While the *PRKAG1* and *PRKAG2* probes showed a broad tissue distribution of expression, *PRKAG3* showed a distinct muscle-specific expression. This result is also supported by the human EST database where multiple ESTs  
30 representing *PRKAG1* and *PRKAG2* have been identified in various cDNA libraries whereas a single EST (GenBank entry AA178898) representing *PRKAG3* has been obtained from a muscle cDNA library. The muscle-specific expression of *PRKAG3* and the lack of expression in liver  
35 are entirely consistent with the phenotypic effect of RN, namely that glycogen content is altered in muscle but

normal in liver (ESTRADE et al., Comp. Biochem. Physiol. 104B, 321-326, 1993).

*PRKAG3* sequences were determined from *rn<sup>+</sup>/rn<sup>+</sup>* and *RN<sup>-</sup>/RN<sup>-</sup>* homozygotes by RT-PCR analysis. A comparison  
5 revealed a total of seven nucleotide differences four of which were nonsynonymous substitutions was found between the sequence from *rn<sup>+</sup>* and *RN<sup>-</sup>* animals, as shown in Table 3 below. Screening of these seven SNPs with genomic DNA from additional *rn<sup>+</sup>* and *RN<sup>-</sup>* pigs of different breeds  
10 revealed five different *PRKAG3* alleles, but only the R41Q missense substitution was exclusively associated with *RN<sup>-</sup>*. This nonconservative substitution occurs in CBS1 which is the most conserved region among isotypic forms of the AMPK  $\gamma$  chain and arginine at this residue (number 70 in  
15 *Prkag1*) is conserved among different isoforms of mammalian AMPK  $\gamma$  sequences as well as in the corresponding *Drosophila* sequence (Figure 3). A simple diagnostic DNA test for the R41Q mutation was designed based on the oligonucleotide ligation assay (OLA; LANDEGREN et al.,  
20 Science, 241, 1077-1080, 1988). Screening a large number of *RN<sup>-</sup>* and *rn<sup>+</sup>* animals from the Hampshire breed as well as large number of *rn<sup>+</sup>* animals from other breeds showed that the 41Q allele was present in all *RN<sup>-</sup>* animals but not found in any *rn<sup>+</sup>* animals, as shown in Table 4 below. The  
25 absence of the 41Q allele from other breeds is consistent with the assumption that the *RN<sup>-</sup>* allele originated in the Hampshire breed; the allele has not yet been found in purebred animals from other breeds. In conclusion, the results provide convincing evidence that *PRKAG3* is  
30 identical to the *RN* gene and that the R41Q substitution most likely is the causative mutation.

Table 3. Comparison of the *PRKAG3* sequences associated with the *rn*<sup>+</sup> and *rn*<sup>-</sup> alleles in different pig populations<sup>a</sup>

Associated allele	RN allele	nt83	nt152	Codon							Population <sup>b</sup>
				34	35	40	41	213			
	RN <sup>-</sup>	ACC	CTC	GCC	CTG	GTC	CAA	TCT		H	
		T	L	A	L	V	Q	S			
	rn <sup>+</sup>	---	---	---	---	---	-G-	---		L, LW, WB	
		-	-	-	-	-	R	-			
	rn <sup>+</sup>	---	-C-	--T	T--	---	-G-	--C		H, L, LW, M, WB	
		-	P	-	-	-	R	-			
	rn <sup>+</sup>	-A-	-C-	--T	T--	---	-G-	--C		D, H	
		N	P	-	-	-	R	-			
	rn <sup>+</sup>	---	-C-	--T	T--	A--	-G-	--C		H, LW, WB, D, L	
		-	P	-	-	I	R	-			

nucleotide and codon numbers refer to the numbering of the sequence SEQ ID NO: 1  
 I=Hampshire, L=Landrace, LW=Large White, M=Meishan, WB=Wild Boar, D=Duroc  
 .D.=not determined, "--" indicates identity to the top sequence.

TABLE 4

RN phenotype	Genotype at nucleotide 593 <sup>a</sup>			Total
	A/A	G/A	G/G	
RN <sup>-</sup> , Hampshire <sup>a</sup>	40	87	0	127
RN <sup>-</sup> , Hampshire <sup>a,b</sup>	0	13	0	13
m <sup>+</sup> , Hampshire <sup>a</sup>	0	0	60	60
rn <sup>+</sup> , other breeds <sup>c</sup>	0	0	488	488

<sup>a</sup>represent both French and Swedish Hampshire populations

<sup>b</sup>heterozygosity RN/m<sup>+</sup> deduced using pedigree information

5 <sup>c</sup>breeds: Angler Saddleback, n=31; Blond Mangalitza, n=2; Bunte Bentheimer, n=16; Duroc, n=160; Göttinger Minipig, n=4; Landrace, n=83; Large White, n=72; Meishan, n=8; Piétrain, n=75; Red Mangalitza, n=5; Rotbunte Husumer, n=15; Schwalbenbauch Mangalitza, n=7; Schwäbisch Hällische, n=2; European Wild Boar, n=5; Japanese Wild Boar, n=3.

10 <sup>d</sup>refers to the nucleotide numbers of SEQ ID NO: 1

Without being bound to any particular mechanism, it may be hypothesised that the AMPK heterotrimer including PRKAG3 is involved in the regulation of glucose transport into skeletal muscle.

15 It has recently been reported that AMPK activation induced by the AMP analogue AICAR or by muscle contraction leads to an increased glucose uptake in skeletal muscle (BERGERON et al., Am. J. Physiol., 276, E938-944, 1999; HAYASHI et al., Diabetes, 47, 1369-1373, 20 1998). If this is the function of the AMPK heterotrimer including PRKAG3, R41Q may be a gain-of-function mutation causing a constitutively active holoenzyme, for instance due to the loss of an inactivating allosteric site. If so, the reduced AMPK activity in RN<sup>-</sup> animals is likely to 25 reflect feed-back inhibition due to the high-energy status of the muscle. An increased uptake of glucose to skeletal muscle is expected to lead to an increase in muscle glycogen content as observed in RN<sup>-</sup> animals. It has been shown that overexpression of glucose transporter 4 30 (GLUT4) in transgenic mice leads to increased uptake of glucose and increased glycogen storage (TREADWAY et al., J. Biol. Chem., 269, 29956-29961, 1994). This type of gain-of-function model is consistent with the dominance

of RN<sup>-</sup> as the presence of a single unregulated copy would have a large effect on AMPK enzyme activity.

An alternative hypothesis on the functional significance of the R41Q substitution associated with the RN<sup>-</sup> allele may also be proposed. Based on the established roles of the yeast SNF1 enzyme in utilisation of glycogen and of mammalian AMPK for inhibiting energy-consuming pathways and stimulating energy-producing pathways, activated AMPK is expected to inhibit glycogen synthesis and stimulate glycogen degradation. If this is the functional role of the isoform(s) containing the PRKAG3 product, the R41Q substitution would be a loss-of-function mutation or a dominant-negative mutation locking the AMPK heterotrimer in an inactive state, and thus inhibiting AMP activation and glycogen degradation. In these cases the phenotypic effect should be explained by haplo-insufficiency, since RN<sup>-</sup> appears fully dominant.

R41Q may thus be a dominant negative mutation, but only if it interferes with multiple isoforms since the major AMPK activity in muscle appears to be associated with the PRKAG1 and 2 isoforms [CHEUNG, et al. *Biochem. J.* 346, 659 (2000)].

The distinct phenotype of the RN<sup>-</sup> mutation indicates that PRKAG3 plays a key role in the regulation of energy metabolism in skeletal muscle. For instance, PRKAG3 is likely to be involved in the adaptation to physical exercise, which is associated with increased glycogen storage. It is also conceivable that loss-of-function mutations in PRKAG3 (or other AMPK genes) may predispose individuals to noninsulin-dependent diabetes mellitus, and AMPK isoforms are potential drug targets for treatment of this disorder.

#### EXAMPLE 2: DETECTION OF THE R41Q SUBSTITUTION IN PIG PRKAG3

A part of PRKAG3 including codon 41 was amplified in 10 µl reactions containing 100 ng genomic



DNA, 0.2 mM dNTPs, 1.5 mM MgCl<sub>2</sub>, 4.0 pmol of both forward (AMPKG3F3:5'-GGAGCAAATGTGCAGACAAG-3') and reverse (AMPKG3R2:5'-CCCACGAAGCTCTGCTTCTT-3') primer, 10% DMSO, 1 U of Taq DNA polymerase and reaction buffer (ADVANCED BIOTECH, London, UK). The cycling conditions included an initial incubation at 94°C for 5 min followed by 3 cycles at 94°C (1 min), 57°C (1 min) and 72°C (1 min), and 35 cycles of 94°C (20 sec), 55°C (30 sec) and 72°C (30 sec). Allele discrimination at nucleotide position 122 was done using the oligonucleotide ligation assay (OLA, LANDEGREN *et al.*, Science, 241, 1077-1080, 1988). The OLA method was carried out as a gel-based assay. Each 10 µl OLA reaction contained 0.5 pmol of each probe SNPRN-A (5'Hex-TGGCCAACGGCGTCCA-3'), SNPRN-G (5'ROX-GGCCAACGGCGTCCG-3') and SNPRN-Common (5'phosphate-AGCGGCACCTTTGTGAAAAAAAAA-3'), 1.5 U of thermostable AMPLIGASE and reaction buffer (EPICENTRE TECHNOLOGIES, Madison, WI) and 0.5 µl of the AMPKG3F3/AMPKG3R2 PCR product. After an initial incubation at 95°C for 5 min, the following thermocycling profile was repeated 10 times: denaturation at 94°C (30 sec), and probe annealing and ligation at 55°C (90 sec). After OLA cycling, 1 µl of product was heat denatured at 94°C (3 min), cooled on ice, and loaded onto 6% polyacrylamide denaturing gel for electrophoresis on an ABI377 DNA sequencer (PERKIN ELMER, Foster City, USA). The resulting fragment lengths and peak fluorescence were analysed using GENESCAN software (PERKIN ELMER, Foster City, USA).

The OLA-based method for the R41Q mutation was used to determine the genotype of DNA samples collected from 68 Swedish Hampshire animals phenotyped as either RN<sup>-</sup> or rn<sup>+</sup> based on their glycolytic potential (GP) value. Figure 6 illustrates typical OLA results from the three possible genotypes. All RN<sup>-</sup> animals were scored as homozygous A/A (n=28) or heterozygous A/G (n=36) at

nucleotide position 122 whereas the  $rn^+$  animals were homozygous G/G (n=4) at this position.

**EXAMPLE 3: PREDICTING THE PRESENCE OF THE  $RN^-$  ALLELE USING A CLOSELY LINKED MICROSATELLITE, MS127B1**

5 A microsatellite 127B1 (MS127B1) was cloned from BAC 127G7 containing pig *PRKAG3*. The BAC clone was digested with *Sau3AI* and the restriction fragments subcloned into the *BamHI* site of pUC18. The resulting library was probed with a (CA)<sub>15</sub> oligonucleotide probe labelled with [ $\gamma$ -<sup>32</sup>P]-  
10 dATP. Strongly hybridising clones were sequenced and primers for PCR amplification of microsatellite loci were designed. Ten  $\mu$ l PCR reactions were performed containing 100 ng genomic DNA, 0.2 mM dNTPs, 1.5 mM MgCl<sub>2</sub>, 4.0 pmol of both forward (MS127B1F:5'-Fluorescein-  
15 CAAACTCTTCTAGGCGTGT-3') and reverse (MS127B1R:5'-GTTTCTGGAACTTCCATATGCCATGG-3') primers, and 1 U of *Taq* DNA polymerase and reaction buffer (ADVANCED BIOTECH, London, UK). The cycling conditions included an initial incubation at 94°C for 5 min followed by 3 cycles at 94°C  
20 (1 min), 57°C (1 min) and 72°C (1 min), and 35 cycles of 94°C (20 sec), 55°C (30 sec) and 72°C (30 sec). The PCR products (0.3  $\mu$ l) were separated using 4% polyacrylamide denaturing gel electrophoresis on an ABI377 DNA sequencer (PERKIN ELMER, Foster City, USA). The resulting fragment  
25 lengths were analysed using the GENESCAN and GENOTYPER software (PERKIN ELMER, Foster City, USA).

The method was used to determine the genotype of DNA samples collected from 87 Swedish Hampshire animals phenotyped as either  $RN^-$  or  $rn^+$  based on their  
30 glycolytic potential (GP) value. Allele 108 (bp) showed a complete association to the  $RN^-$  allele in this material as all  $RN^-$  ( $RN^-/RN^-$  or  $RN^-/rn^+$ ) animals were homozygous or heterozygous for this allele while no  $rn^+$  ( $rn^+/rn^+$ ) animals carried this allele, as shown in Table 5 below.

TABLE 5

Animals	n	Genotype				
		94/94	94/108	94/114	100/108	108/108
RN <sup>-</sup>	80	0	37	0	2	41
rn <sup>+</sup>	7	3	0	4	0	0

**EXAMPLE 4: DETECTING THE PRESENCE OF THE RN<sup>-</sup> ALLELE USING A PCR-RFLP TEST**

The RN<sup>-</sup> mutation inactivates a BsrBI site  
 5 GAG<sup>^</sup>CGG/CTC<sup>^</sup>GCC (BsrBI RE site is not palindromic). At that site, the RN<sup>-</sup> sequence is AAGCGG instead of GAGCGG.

A 134 bp long fragment of the RN gene is amplified from porcine genomic DNA. The rn<sup>+</sup> allele is identified after BsrBI digestion, by detection of two  
 10 fragments of 83 and 51 bps.

The test is performed as follows:

1° Primer sequences:

Sequence of primers used to amplify the RN mutation region:

15 RNU: 5' GGGAACGATTCACCCTCAAC 3'  
 RNL: 5' AGCCCCTCCTCACCCACGAA 3'

To provide an internal control of digestion, a BsrBI site has been added at the extremity of one of the two primers within a 20 bp long tail. The tail permits  
 20 both creation of a BsrBI site (a shorter tail might be sufficient), and an easy discrimination of uncut fragment from other fragments. The use of tailed primers does not affect efficiency and specificity of amplification.

The sequence of the RNL modified primer  
 25 including a control tail with a BsrBI site is:

RNLBsrA14: 5'

A<sub>5</sub>C<sub>2</sub>A<sub>7</sub>CCGCTCAGCCCCTCCTCACCCACGAA 3'

2° PCR reaction mixture used:

50 ng DNA  
 0.5 Unit Taq polymerase (GIBCO BRL)  
 1.5 mM MgCl<sup>2</sup>  
 200 mM dNTP  
 30

0.2  $\mu$ M each primer

Total reaction volume: 25  $\mu$ l

3° PCR conditions used (on OMNIGENE HYBAID thermocycler):

- 5                    1x (5min 95°C)  
                     35x (45sec 57°C, 45sec 72°C, 45sec 95°C)  
                     1x (45sec 57°C, 15min 72°C)

4° Restriction enzyme digestion performed at 37°C for 2 hours:

- 10                   10  $\mu$ l PCR product  
                     1x BsrBI BIOLABS buffer  
                     5U BsrBI restriction enzyme (BIOLABS)  
                     Total reaction volume: 15  $\mu$ l

5° Size of fragments produced after PCR using primers with control tail and digestion with BsrBI:

- 15                   Uncut fragment from *RN* or *rn*<sup>+</sup> allele : 154 bp  
                     After digestion of fragment amplified from *RN*  
                     allele : 137 bp + 17 bp  
                     After digestion of fragment amplified from *rn*<sup>+</sup>  
                     allele : 83 bp + 54 bp + 17 bp  
                     Size difference can be identified either after  
                     polyacrylamide, agarose/NUSIEVE or agarose gel  
                     electrophoresis.

**EXAMPLE 5: EFFECT OF V40I POLYMORPHISM ON GLYCOLYTIC POTENTIAL.**

25                   Further, a set of 181 *rn*<sup>+</sup>/*rn*<sup>+</sup> homozygous  
                     animals (R/R at position 41 of SEQ ID NO: 2) were  
                     analyzed for the V40I polymorphism (referring to position  
                     40 of SEQ ID NO: 2) by PCR-RFLP using *Fok*I restriction  
                     enzyme. The glycolytic potential was determined in  
                     parallel according to the method disclosed by MONIN et  
                     al., (Meat Science, 13, 49-63, 1985).

The results are shown in Table 6 below:

Table 6

Genotype at position 40	Average glycolytic potential	Standard Deviation	Number of typed animals
I/I	178.30	31.13	13
V/I	204.15	37.73	164
V/V	210.83	38.21	104

These results show that the V40I polymorphism has a significant effect on the glycolytic potential in skeletal muscle.

## CLAIMS

- 1) A gamma subunit of a vertebrate AMP-activated kinase (AMPK), wherein said gamma subunit is a polypeptide comprising at least a sequence having at least 70% identity with the polypeptide SEQ ID NO: 2.
- 2) A polypeptide of claim 1, wherein said polypeptide comprises a sequence having at least 95% identity with the polypeptide SEQ ID NO:2.
- 3) A polypeptide of claim 1, wherein said polypeptide comprises a sequence having at least 75% identity with the polypeptide SEQ ID NO: 28.
- 4) A polypeptide of any of claims 1 to 3, wherein said polypeptide comprises the sequence SEQ ID NO: 2 or SEQ ID NO:4.
- 5) A polypeptide of claim 4, wherein said polypeptide comprises the sequence SEQ ID NO: 28, SEQ ID NO: 30 or SEQ ID NO: 32.
- 6) A polypeptide which is a functionally altered mutant of a gamma subunit of a vertebrate AMP-activated kinase, wherein said polypeptide has at least a mutation located within the first CBS domain of said gamma subunit.
- 7) A polypeptide of claim 6, wherein the mutation is located within the region of the first CBS domain aligned with the region of a polypeptide of SEQ ID NO: 2 spanning from residue 30 to residue 50.
- 8) A polypeptide of claim 7, wherein the mutation is a R→Q substitution or a V→I substitution.
- 9) A polypeptide of claim 8 selected among:
- a polypeptide having a sequence resulting from a R→Q substitution at a position corresponding to position 41 in SEQ ID NO: 2;
  - a polypeptide having a sequence resulting from a V→I substitution at the position corresponding to position 40 of SEQ ID NO: 2.

10) A polypeptide which is a mutant of a gamma subunit of a vertebrate AMP-activated kinase, wherein said polypeptide results from a deletion of a part of a polypeptide of any of claims 1 to 5.

5 11) A nucleic acid sequence encoding a polypeptide of any of claims 1 to 10, or the complement thereof, provided that said nucleic acid sequence does not consist of the EST GENBANK AA178898, or of the EST W94830.

10 12) A nucleic acid sequence of claim 11, having the sequence SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, or the complement thereof.

15 13) A nucleic acid sequence comprising at least a portion of a nucleic acid sequence encoding a polypeptide of any of claims 1 to 10, and up to 500 kb of a 3' and/or of a 5' adjacent genomic DNA sequence, or the complement thereof.

20 14) A nucleic acid fragment selected among:  
- a specific fragment of a nucleic acid sequence encoding a polypeptide of any of claims 1 to 10, or of a nucleic acid sequence of claim 13;  
- a nucleic acid fragment which specifically hybridises under stringent conditions with a nucleic acid sequence  
25 encoding a polypeptide of any of claims 1 to 8, or of a nucleic acid sequence of claim 11;  
provided that said nucleic acid fragment does not consist of the EST GENBANK AA178898 or of the EST GENBANK W94830.

30 15) A set of primers for amplifying a nucleic acid sequence of any of claims 11 to 13 or a portion thereof, comprising at least a primer consisting of a nucleic acid fragment of claim 14.

35 16) A recombinant vector comprising a nucleic acid sequence encoding a polypeptide of any of claims 1 to 10.

17) An host cell transformed by a nucleic acid sequence encoding a polypeptide of any of claims 1 to 10.

18) A transgenic animal transformed by a nucleic acid sequence encoding a polypeptide of any of  
5 claims 1 to 10.

19) A knockout animal, wherein the gene encoding a polypeptide of any of claims 1 to 5 is inactive.

20) A heterotrimeric AMPK wherein the  $\gamma$  subunit consists of a polypeptide of any of claims 1 to  
10 10.

21) A method of detecting a metabolic disorder resulting from a mutation in a gene encoding a  $\gamma$  subunit of AMPK, wherein said process comprises:

15 - obtaining a nucleic acid sample from a vertebrate;

- checking the presence in said nucleic acid of a nucleic acid sequence encoding a polypeptide of any of claims 1 to 10, wherein said polypeptide is  
20 functionally altered.

22) A method of claim 21 wherein the disorder is correlated with an altered glycogen accumulation in the muscular cells and results from the expression of a functionally altered allele of a polypeptide of any of  
25 claims 1 to 5.

23) A method of any of claims 21 or 22 wherein the presence of the nucleic acid sequence encoding said mutant polypeptide is checked by contacting said nucleic acid sample with a nucleic acid probe obtained from a  
30 nucleic acid of claim 14 and spanning said mutation, under conditions of specific hybridisation between said probe and the mutant sequence to be detected, and detecting the hybridisation complex..

24) A method for obtaining a pair of primers  
35 allowing to detect a genetic polymorphic marker linked to



a nucleic acid sequence encoding a polypeptide of any of claims 1 to 5, wherein said process comprises:

- screening a genomic DNA library from a vertebrate with a probe specific for a nucleic acid sequence encoding a polypeptide of any of claims 1 to 5, in order to select clones comprising said nucleic acid sequence and flanking chromosomal sequences;
- identifying a polymorphic locus in said flanking chromosomal sequences, and sequencing a DNA segment comprising said polymorphic locus ;
- designing primer pairs flanking said polymorphic locus.

25) A method of claim 24 wherein the selected clones comprise at least a portion of a nucleic acid sequence encoding a polypeptide of any of claims 1 to 5, and up to 500 kb of a 3' and/or of a 5' adjacent sequence.

26) A method of any of claims 21 to 25 wherein the vertebrate is a mammal.

27) A method of claim 26 wherein said mammal is a pig.

28) A pair of primers obtainable by the process of any of claims 24 to 26.

29) A process for detecting a dysfunction of carbohydrate metabolism resulting from the expression of a functionally altered allele of a polypeptide of any of claims 1 to 5 in a vertebrate, wherein said process comprises:

- obtaining a sample of genomic DNA from said vertebrate;
- contacting said DNA with a pair of primers of claim 28 under conditions allowing PCR amplification;
- analysing the PCR product to detect if an allele of a polymorphic marker linked to a nucleic acid sequence encoding a functionally altered allele of a polypeptide of any of claims 1 to 5 is present.

30) A process of claim 29, wherein said functionally altered polypeptide results from a R41Q substitution in SEQ ID NO: 2.

31) A process of any of claims 29 or 30,  
5 wherein said vertebrate is a mammal.

32) A process of claim 31 wherein said mammal is a pig.

33) A process of claim 32 wherein the pair of primers is selected among:

10 - a pair of primers consisting of SEQ ID NO: 5 and SEQ ID NO: 6;

- a pair of primers consisting of SEQ ID NO: 7 and SEQ ID NO: 8;

15 - a pair of primers consisting of SEQ ID NO: 9 and SEQ ID NO: 10;

- a pair of primers consisting of SEQ ID NO: 11 and SEQ ID NO: 12;

- a pair of primers consisting of SEQ ID NO: 13 and SEQ ID NO: 14;

20 - a pair of primers consisting of SEQ ID NO: 15 and SEQ ID NO: 16;

- a pair of primers consisting of SEQ ID NO: 17 and SEQ ID NO: 18;

25 - a pair of primers consisting of SEQ ID NO: 19 and SEQ ID NO: 20;

- a pair of primers consisting of SEQ ID NO: 21 and SEQ ID NO: 22;

- a pair of primers consisting of SEQ ID NO: 23 and SEQ ID NO: 24;

30 - a pair of primers consisting of SEQ ID NO: 25 and SEQ ID NO: 26.

34) Use of a transformed cell of claim 17 to screen compounds able to modulate AMPK activity.

35) Use of a transgenic animal of claim 18 to  
35 screen compounds able to modulate AMPK activity.

36) Use of a knockout animal of claim 19 to screen compounds able to modulate energy metabolism in the absence of a functional polypeptide of any of claims 1 to 5.

5           37) Use of an heterotrimeric AMPK of claim 20 to screen compounds able to modulate AMPK activity.

1/8




Figure 1

5'UTR 10 20 30 40 50 60 70 80  
 Pig TTCCTAGAGCAAGGAGAGAGCCGTTTCATGGCCATCCCGAGCTGTAACCACCAGCTCAGAAAGAAGCCATGGGGACCAGGG  
 Hum -----A-A-C---A-C-----A-C-----G-----G-----T--G---A-A-G-A-  
 90 100 110 120 130 140 150 160  
 Pig GAACAAGGCCTCTAGATGGACAAGGCAGGAGGATGTAGAGGAAGGGGGCCTCCGGGCCCGAGGGAAGGTCCCCAGTCCA  
 Hum -GC---A---TG-----A---TCG--G-----A---A---A---T-A-G-----G-----  
 170 180 190 200 210 220 230 240  
 Pig GGCCAGTTGCTGAGTCCACCGGGCAGGAGGCCACATTCCCCAAGGCCACACCCTTGGCCCAAGCCGCTCCCTTGCCCGAG  
 Hum -----AC-----T-----A-----T-----T-A-----T-----G-----  
 250 260 270 280 290 300 310 320  
 Pig GTGGACAACCCCCCAACAGAGCGGGACATCCTCCCTCTGACTGTGCAGCCTCAGCCTCGACTCCAACACAGACCATCT  
 Hum ----G--CT--A-----G-T---TG-----A-----TG-A-G-----G-----TG--G-  
 330 340 350 360 370 380 390 400  
 Pig GGATCTGGGCATAGAGTTCTCAGCCTCGGCGCGTCCGGGGATGAGCTTG...GGCTGGTGGAAGAGAAGCCAGCCCCGT  
 Hum ---G---C--CG-----C-----A-A-A--C-G--A-TG-----A-AAG-C---C-----G---T---T---  
 410 420 430 440 450 460 470 5'UTR  
 Pig GCCCATCCCCAGAGGTGCTGTTACCCAGGCTGGGCTGGGATGATGAGCTGCAGAAGCCGGGGGCCAGGTCTAC  
 Hum ---TG-----GC---CC-CA--T---A-----C--A---G---A--C--C-----A-----  
 CDS 10 20  
 Pig ATG CAC TTC ATG CAG GAG CAC ACC TGC TAC GAT GCC ATG GCG ACC AGC TCC AAA CTG GTC  
 . Met His Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser Lys Leu Val  
 Hum --- -G- --- --- --- --- --- --- --- --- ---A---T--- --- ---G---A---  
 - Arg -  
 30 40  
 Pig ATC TTC GAC ACC ATG CTG GAG ATC AAG GCC TTC TTT GCC CTG GTG GCC AAC GGC GTC  
 Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe Ala Leu Val Ala Asn Gly Val  
 Hum --- --- --- --- --- --- --- --- ---T--- --- --- ---T---G---  
 50 60  
 Pig CGA GCG GCA CCT TTG TGG GAC AGC AAG AAG CAG AGC TTC GTG GGG ATG CTG ACC ATC ACA  
 Arg Ala Ala Pro Leu Trp Asp Ser Lys Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr  
 Hum --G--A--C---C-A--- --- --- ---T--- --- --- --- ---T---  
 70 80  
 Pig GAC TTC ATC TTG GTG CTG CAC CGC TAT TAC AGG TCC CCC CTG GTC CAG ATC TAC GAG ATT  
 Asp Phe Ile Leu Val Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile  
 Hum --- --- --- C--- --- ---T--- ---C--- --- --- ---T--- ---  
 90 100  
 Pig GAA GAA CAT AAG ATT GAG ACC TGG AGG GAG ATC TAC CTT CAA GGC TGC TTC AAG CCT CTG  
 Glu Glu His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe Lys Pro Leu  
 Hum --- C--- --- --- --- --- --- ---G--- --- --- --- ---  
 - Gln -  
 110 120  
 Pig GTC TCC ATC TCT CCC AAT GAC AGC CTG TTC GAA GCT GTC TAC GCC CTC ATC AAG AAC CGG  
 Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val Tyr Ala Leu Ile Lys Asn Arg  
 Hum --- --- --- ---T--- ---T--- ---T--- ---A--- --- --- --- ---  
 -  
 130 140  
 Pig ATC CAC CGC CTG CCG GTC CTG GAC CCT GTC TCC GGG GCT GTG CTC CAC ATC CTC ACA CAT  
 Ile His Arg Leu Pro Val Leu Asp Pro Val Ser Gly Ala Val Leu His Ile Leu Thr His  
 Hum --- --T--- --- --T---T---T--- --G--G--A--C AAC --A--- --- --- ---C---  
 -  
 150 160  
 Pig AAG CCG CTT CTC AAG TTC CTG CAC ATC TTT GGC ACC CTG CTG CCC CGG CCC TCC TTC CTC  
 Lys Arg Leu Leu Lys Phe Leu His Ile Phe Gly Thr Leu Leu Pro Arg Pro Ser Phe Leu  
 Hum --A--C--G--- --- --- ---T--- --- --- --- --- --- --- ---  
 -  
 Ser -

Figure 2

**Figure 2 (cont.)**

4/8

```

      410      420      430      440      450      460      470      480
Pig GGGAGTTAGAGCTGCCCTCTCTCAGTT.....CAGTTCCTCCCTGCTGAGAATG.TCCCTGGAAGGAAGCCAGTTAAT
Hum -----CCC-----TTG-.-C---CAACGTGCG--C-G---T---A---CTCC-G-C-TTG-CATTTTC---G-T-C-G-.
      490      500      510      520      530      540      550      560
Pig AAACCTTGGTTGGATGGAATTTCCCACTCG.....
Hum --TG--GCA--TC-G--G.....CA-G-AGCAGCCGTTATTTATAGAACTGCCTGTTGGAGGTGGGAGTCCCTCCCT
      570      580      590      600      610      620      630      640
Pig .....
Hum CCATTCTTGTCCAGAAACTCCTTAGCTCTCGCAGTGAGCCATGTTCTTAGTCTCCAGGGATGGATGGCCTTGTATATGC
      650      660      670      680      690      700      710      720
Pig .....
Hum ACCCCTGAGAATGAGCAATTGAGAAAACAAAACAAAAGGAACAATCCATGAACCTTAGATTATTATTGTTTCACTCAAAAT

      730      740
Pig .....
Hum GCTGCAGTCATTTGACCTG
```

Figure 2 (cont.)

PigG3 MSFLEQGESRSHPSRAVTTSSERSHGDQGNKASRWTRQEDVEEGPPGREGPQSRPVAESTGQEA TP KATPLAQAPLAEVNDNPTERDILPSDCAAS 100  
HumG3 -----EN-S-----P-----S-----IR-KRRA--L-----KS-----E-----QQ--R--T-----L-----D--G-GT--GW-C-----T--  
CB91→ 200

PigG3 ASDNTOHLDLGIEFSASASGDEL.GLVEEKPA PCSPREVLLPRLGWDELQKPGAQVYMIHQEHTCYDAMATSSKLVIPTMLEIKKAFFALVANGV 300  
HumG3 -AG-S--DVE-AT--P-TE-WEC--E--L--R--L--L--QAPP-K-----R-----I--R-----V--S-QV-----T--  
HumG1 METVIS-DSSPAVENEHPQ-TPESNNS--TS--KS-R--LIP-----V--S-QV-----T--  
HumG2 AALGPAREGM-EKLEFE-EAVEDSESG--R--RS-K--IVP-----V--T-QV-----T--  
Dros RDSRGLPVADPPFLEKVNLS-D-EEDDS-IFVK-FRP-K--LIP-A--V--Q-LV--Y--Y--  
Snf4 MK-TQDSQEKVSEIQQLAVES..IRK-LNSK-S--VLPV-YR-IVL--S-LV--SLNV-LQ-SI  
CB92→ 300

PigG3 KAAPLRDSKKQSPVGMCTTDPILVLRHYRSPVQIYEIEENKIETWREIYLQCFKPL..VSTSPNDSLFVAVYALTKNRIHRLPVDPSCA...V 400  
HumG3 -----Q-----V--DS-----C--A--D--SS-R-K-----I--E--N--T--  
HumG1 -----NI--K-A--L-----L-----V--DS-----C--A--D--SS-R-K-----I--E--N--T--  
HumG2 -----E--T--SL-----NI--K-M--L-----L-----ET-----N--DA--D--S--K-----I--I--N--A--  
Dros -----E--Q-----KI-QM--K--NASMEQL--LD--DV--HNQVM--G-DA--YD-IKI--HS-----I--AT--N--  
Snf4 VS-----TSR-A-L--T--N-IQY-FSN-D.KPELVOKLQDLGLKD-ERALGVQD-Q-DTA--H-SRP--CLKNLES-SG-I-LI-QDEETUREI-  
CB93→ 400


PigG3 LHILTHIKRLKFLHIFGTLPRPSPFLYRTIQDLGIGTFPRDLAVVLETA PILTALDIFVDRRVSA LPVNETQGVGLYSRFDV IHLAAQQTYNHILDMNVG 400  
HumG3 -----S-----Y--I--KL-I-EP-K-E-MKSLEE-Q--YANI-M-RT-T-VVV-G--QH-----D-K-R-DI-K--N--EK--N--VS-T--  
HumG1 -----I--I--QL-MSDM-K-A-MKQNLDB-----YINI-FIHPDT--IK-N--E--I-----D-S-K-DI-K--N--EK--N--IT-T--  
HumG2 -----I--R--FLYINE--K-AYMQKSLRE-K--YNNIETAD--TS-I--KK-E-----L-DSD-RL-DI-AK--N--EK--D--VSLR--  
Dros VSV--QY-I--VALNCRE...TH--KIP-G--N-I-QDNMKSCQM-T-VIDVIQMLTQG--SV-IID-N-YLINV-EAY--LG-IKGGI--D-SLS--  
Snf4 CB94→ 472

PigG3 EALRQRTLCLEQVLSQCPHETLGEVIDRIVREQVHRLVLDVETQHLLGVVSLSDILQALVLS PAGIDALGA\* 472  
HumG3 -----S-----K--Y-L-----ETI-N-L-EAE-----V--NDVVK-I-----TGGEKKP\*  
HumG1 K--QH-SHYP-----K--Y-L-----ETI-N-L-EAE-----V--NDVVK-I-----TGGEKKP\*  
HumG2 Q--QH-SQYF-----VK-NKL-I-ETIV-----AE-----V-N-ADSV-II-----I-T--AKQKETETE\*  
Dros K-NEH-NEWP-----QK-NLD-S-YTIME-----AE-----V--NRKVI-II-----LY--R-S-EGV  
Snf4 ---MR-SDDP-----YT-TKNDK-STIM-N-RKAR---FFV--DVGR-V--LT-----KYIL-GSN\*

Figure 3



6/8

**Figure 4**





Figure 5

8/8

**Figure 6**

1/20

## SEQUENCE LISTING

<110> INSTITUT NATIONAL DE LA RECHERCHE AGRONOMIQUE  
 MILAN, Denis  
 ANDERSSON, Leif  
 LOOFT, Christian  
 ROBIC, Annie  
 ROGEL-GAILLARD, Claire  
 IANNUCELLI, Nathalie  
 GELLIN, Joël  
 KALM, Ernst  
 LE ROY, Pascale  
 CHARDON, Patrick

<120> VARIANTS OF THE GAMMA CHAIN OF AMPK, DNA SEQUENCES ENCODING  
 THE SAME, AND USES THEREOF

<130> MJPCb539-99

<140>  
 <141>

<150> EP 99402236.3  
 <151> 1999-09-10

<150> EP 00401388.4  
 >151> 2000-05-18

<160> 32

<210> 1  
 <211> 1867  
 <212> DNA  
 <213> Sus scrofa

<220>  
 <221> CDS  
 <222> (472) .. (1389)

<400> 1  
 ttcctagagc aaggagagag ccgttcatgg ccatcccgag ctgtaaccac cagctcagaa 60  
 agaagccatg gggaccaggg gaacaaggcc tctagatgga caaggcagga ggatgtagag 120  
 gaaggggggc ctccgggccc gagggaaggt cccagtgcca ggccagtgc tgagtccacc 180  
 gggcaggagg ccacattccc caaggccaca cccttgccc aagccgctcc cttggccgag 240  
 gtggacaacc cccaacaga gcgggacatc ctccctctg actgtgcagc ctcagcctcc 300  
 gactccaaca cagaccatct ggatctgggc atagagttct cagcctcggc ggcgtcgggg 360  
 gatgagcttg ggctggtgga agagaagcca gccccgtgcc catccccaga ggtgctgtta 420  
 cccaggctgg gctgggatga tgagctgcag aagccggggg cccagggtcta c atg cac 477  
 Met His  
 1

ttc atg cag gag cac acc tgc tac gat gcc atg gcg acc agc tcc aaa 525  
 Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser Lys  
 5 10 15

2/20

ctg gtc atc ttc gac acc atg ctg gag atc aag aag gcc ttc ttt gcc	573
Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe Ala	
20 25 30	
ctg gtg gcc aac ggc gtc cga gcg gca cct ttg tgg gac agc aag aag	621
Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser Lys Lys	
35 40 45 50	
cag agc ttc gtg ggg atg ctg acc atc aca gac ttc atc ttg gtg ctg	669
Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu Val Leu	
55 60 65	
cac cgc tat tac agg tcc ccc ctg gtc cag atc tac gag att gaa gaa	717
His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile Glu Glu	
70 75 80	
cat aag att gag acc tgg agg gag atc tac ctt caa ggc tgc ttc aag	765
His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe Lys	
85 90 95	
cct ctg gtc tcc atc tct ccc aat gac agc ctg ttc gaa gct gtc tac	813
Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val Tyr	
100 105 110	
gcc ctc atc aag aac cgg atc cac cgc ctg ccg gtc ctg gac cct gtc	861
Ala Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp Pro Val	
115 120 125 130	
tcc ggg gct gtg ctc cac atc ctc aca cat aag cgg ctt ctc aag ttc	909
Ser Gly Ala Val Leu His Ile Leu Thr His Lys Arg Leu Leu Lys Phe	
135 140 145	
ctg cac atc ttt ggc acc ctg ctg ccc cgg ccc tcc ttc ctc tac cgc	957
Leu His Ile Phe Gly Thr Leu Leu Pro Arg Pro Ser Phe Leu Tyr Arg	
150 155 160	
acc atc caa gat ttg ggc atc ggc aca ttc cga gac ttg gcc gtg gtg	1005
Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala Val Val	
165 170 175	
ctg gaa acg gcg ccc atc ctg acc gca ctg gac atc ttc gtg gac cgg	1053
Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val Asp Arg	
180 185 190	
cgt gtg tct gcg ctg cct gtg gtc aac gaa act gga cag gta gtg ggc	1101
Arg Val Ser Ala Leu Pro Val Val Asn Glu Thr Gly Gln Val Val Gly	
195 200 205 210	
ctc tac tct cgc ttt gat gtg atc cac ctg gct gcc caa caa aca tac	1149
Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln Thr Tyr	
215 220 225	
aac cac ctg gac atg aat gtg gga gaa gcc ctg agg cag cgg aca ctg	1197
Asn His Leu Asp Met Asn Val Gly Glu Ala Leu Arg Gln Arg Thr Leu	
230 235 240	
tgt ctg gaa ggc gtc ctt tcc tgc cag ccc cac gag acc ttg ggg gaa	1245
Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Thr Leu Gly Glu	
245 250 255	

3/20

```

gtc att gac cgg att gtc cgg gaa cag gtg cac cgc ctg gtg ctc gtg 1293
Val Ile Asp Arg Ile Val Arg Glu Gln Val His Arg Leu Val Leu Val
    260                265                270

gat gag acc cag cac ctt ctg ggc gtg gtg tcc ctc tct gac atc ctt 1341
Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp Ile Leu
    275                280                285                290

cag gct ctg gtg ctc agc cct gct gga att gat gcc ctc ggg gcc tga 1389
Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly Ala
    295                300                305

gaaccttggga acctttgctc tcaggccacc tggcacacct ggaagccagt gaagggagcc 1449

gtggactcag ctctcacttc ccttcagccc cacttgctgg tctggctctt gttcaggtag 1509

gctccgccccg gggccccctgg cctcagcadc agccccctcag tctccctggg caccagatc 1569

tcagactggg gcacctgaa gatgggagtg gccagctta tagctgagca gccttgtagaa 1629

atctaccagc atcaagactc actgtgggac cactgctttg tccattctc agtgaaatg 1689

atggaggggcc tcataagagg ggtggacagg gcctggagta gaggccagat cagtgcgtg 1749

ccttcaggac ctccggggag ttagagctgc cctctctcag ttcagttccc cctgctgag 1809

aatgtccctg gaaggaagcc agttaataaa ccttggttgg atggaatttc cacactcg 1867

<210> 2
<211> 305
<212> PRT
<213> Sus scrofa

<400> 2
Met His Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser
  1                5                10                15

Ser Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe
    20                25                30

Phe Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser
    35                40                45

Lys Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu
    50                55                60

Val Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile
    65                70                75                80

Glu Glu His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys
    85                90                95

Phe Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala
    100                105                110

Val Tyr Ala Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp
    115                120                125

Pro Val Ser Gly Ala Val Leu His Ile Leu Thr His Lys Arg Leu Leu
    130                135                140

```

SUBSTITUTE SHEET (RULE 26)

4/20

Lys Phe Leu His Ile Phe Gly Thr Leu Leu Pro Arg Pro Ser Phe Leu  
 145 150 155 160  
 Tyr Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala  
 165 170 175  
 Val Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val  
 180 185 190  
 Asp Arg Arg Val Ser Ala Leu Pro Val Val Asn Glu Thr Gly Gln Val  
 195 200 205  
 Val Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln  
 210 215 220  
 Thr Tyr Asn His Leu Asp Met Asn Val Gly Glu Ala Leu Arg Gln Arg  
 225 230 235 240  
 Thr Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Thr Leu  
 245 250 255  
 Gly Glu Val Ile Asp Arg Ile Val Arg Glu Gln Val His Arg Leu Val  
 260 265 270  
 Leu Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp  
 275 280 285  
 Ile Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly  
 290 295 300  
 Ala  
 305

<210> 3  
 <211> 2109  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (472)..(1389)

<400> 3  
 ttcctagagc aagaaaacag cagctcatgg ccataccag ctgtgaccag cagctcagaa 60  
 agaatccgtg ggaaacggag ggccaaagcc ttgagatgga caaggcagaa gtcggtggag 120  
 gaaggggagc caccagggtca gggggaaggt ccccggtcca ggccaactgc tgagtccacc 180  
 gggctggagg ccacattccc caagaccaca cccttggtc aagctgatcc tgccgggggtg 240  
 ggcactccac caacaggggtg ggactgcctc cctctgact gtacagcctc agctgcaggc 300  
 tccagcacag atgatgtgga gctggccacg gagttcccag ccacagaggc ctgggagtgt 360  
 gagctagaag gcctgctgga agagaggcct gccctgtgcc tgtccccgca ggccccatct 420  
 cccaagctgg gctgggatga cgaactgcgg aaacccggcg cccagatcta c atg cgc 477

5/20

															Met Arg	
															1	
ttc atg cag gag cac acc tgc tac gat gcc atg gca act agc tcc aag	525															
Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser Lys																
5 10 15																
cta gtc atc ttc gac acc atg ctg gag atc aag aag gcc ttc ttt gct	573															
Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe Ala																
20 25 30																
ctg gtg gcc aac ggt gtg cgg gca gcc cct cta tgg gac agc aag aag	621															
Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser Lys Lys																
35 40 45 50																
cag agc ttt gtg ggg atg ctg acc atc act gac ttc atc ctg gtg ctg	669															
Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu Val Leu																
55 60 65																
cat cgc tac tac agg tcc ccc ctg gtc cag atc tat gag att gaa caa	717															
His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile Glu Gln																
70 75 80																
cat aag att gag acc tgg agg gag atc tac ctg caa ggc tgc ttc aag	765															
His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe Lys																
85 90 95																
cct ctg gtc tcc atc tct cct aat gat agc ctg ttt gaa gct gtc tac	813															
Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val Tyr																
100 105 110																
acc ctc atc aag aac cgg atc cat cgc ctg cct gtt ctt gac ccg gtg	861															
Thr Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp Pro Val																
115 120 125 130																
tca ggc aac gta ctc cac atc ctc aca cac aaa cgc ctg ctc aag ttc	909															
Ser Gly Asn Val Leu His Ile Leu Thr His Lys Arg Leu Leu Lys Phe																
135 140 145																
ctg cac atc ttt ggt tcc ctg ctg ccc cgg ccc tcc ttc ctc tac cgc	957															
Leu His Ile Phe Gly Ser Leu Leu Pro Arg Pro Ser Phe Leu Tyr Arg																
150 155 160																
act atc caa gat ttg ggc atc ggc aca ttc cga gac ttg gct gtg gtg	1005															
Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala Val Val																
165 170 175																
ctg gag aca gca ccc atc ctg act gca ctg gac atc ttt gtg gac cgg	1053															
Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val Asp Arg																
180 185 190																
cgt gtg tct gca ctg cct gtg gtc aac gaa tgt ggt cag gtc gtg ggc	1101															
Arg Val Ser Ala Leu Pro Val Val Asn Glu Cys Gly Gln Val Val Gly																
195 200 205 210																
ctc tat tcc cgc ttt gat gtg att cac ctg gct gcc cag caa acc tac	1149															
Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln Thr Tyr																
215 220 225																



6/20

aac cac ctg gac atg agt gtg gga gaa gcc ctg agg cag agg aca cta 1197  
 Asn His Leu Asp Met Ser Val Gly Glu Ala Leu Arg Gln Arg Thr Leu  
                   230                                  235                                  240

tgt ctg gag gga gtc ctt tcc tgc cag ccc cac gag agc ttg ggg gaa 1245  
 Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Ser Leu Gly Glu  
                   245                                  250                                  255

gtg atc gac agg att gct cgg gag cag gta cac agg ctg gtg cta gtg 1293  
 Val Ile Asp Arg Ile Ala Arg Glu Gln Val His Arg Leu Val Leu Val  
                   260                                  265                                  270

gac gag acc cag cat ctc ttg ggc gtg gtc tcc ctc tcc gac atc ctt 1341  
 Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp Ile Leu  
                   275                                  280                                  285                                  290

cag gca ctg gtg ctc agc cct gct ggc atc gat gcc ctc ggg gcc tga 1389  
 Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly Ala  
                                   295                                  300                                  305

gaagatctga gtcttcaatc ccaagccaac tgcacactgg aagccaatga aggaattgag 1449

aacagcttca tttccccaac cccaatttgc tgggttcagct atgattcagg cttcttcagc 1509

cttccaaaat tgcctttgcc ttacttgtgc tcccagaacc cttegggcat gccagtgca 1569

ccatgggatg atgaaattaa ggagaacagc tgagtcaagc ttggaggtcc ctgaaccaga 1629

ggcactagga ttaccccagg gccatctgtg ctccatgcc gcccatcccc ttgccgctg 1689

actgggtcgg atggccccag tgggtttagt cagggcttct ggattcctcg gtttctgggc 1749

tacctatggc ttcagccttc agctcctggg agtcccagct gttgttccca gcaacgtcgc 1809

cactgccctc ctactctcca ggctttgtca tttcaaggct gctgaaatgc tgcatttcag 1869

gggccaccat ggagcagccg ttatttatag aactgcctgt tggaggtggg gagtcctccc 1929

tccattcttg tccagaaaac tccttagctc tcgcagtgag ccatgttctt agtctccagg 1989

gatggatggc cttgtatatg gaccctgag aatgagcaat tgagaaaaca aaacaaaagg 2049

aacaatccat gaacttagat tttattggtt tcaactcaaaa tgctgcagtc atttgacctg 2109

<210> 4

<211> 305

<212> PRT

<213> Homo sapiens

<400> 4

Met Arg Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser  
           1                                  5                                  10                                  15

Ser Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe  
                   20                                  25                                  30

Phe Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser  
           35                                  40                                  45

7/20

Lys Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu  
 50 55 60  
 Val Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile  
 65 70 75 80  
 Glu Gln His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys  
 85 90 95  
 Phe Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala  
 100 105 110  
 Val Tyr Thr Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp  
 115 120 125  
 Pro Val Ser Gly Asn Val Leu His Ile Leu Thr His Lys Arg Leu Leu  
 130 135 140  
 Lys Phe Leu His Ile Phe Gly Ser Leu Leu Pro Arg Pro Ser Phe Leu  
 145 150 155 160  
 Tyr Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala  
 165 170 175  
 Val Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val  
 180 185 190  
 Asp Arg Arg Val Ser Ala Leu Pro Val Val Asn Glu Cys Gly Gln Val  
 195 200 205  
 Val Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln  
 210 215 220  
 Thr Tyr Asn His Leu Asp Met Ser Val Gly Glu Ala Leu Arg Gln Arg  
 225 230 235 240  
 Thr Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Ser Leu  
 245 250 255  
 Gly Glu Val Ile Asp Arg Ile Ala Arg Glu Gln Val His Arg Leu Val  
 260 265 270  
 Leu Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp  
 275 280 285  
 Ile Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly  
 290 295 300  
 Ala  
 305

<210> 5  
 <211> 20  
 <212> DNA  
 <213> Sus scrofa

<400> 5  
 ggaatttcaa gtcagccaac

8/20

<210> 6  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 6  
cttcaaaaga ccgtgctact 20

<210> 7  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 7  
ctgggaacct ctatatgctg 20

<210> 8  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 8  
tagggaaata caaatcacag 20

<210> 9  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 9  
ctccagctca caggatgaca 20

<210> 10  
<211> 26  
<212> DNA  
<213> Sus scrofa

<400> 10  
gtttctgcag ctttagcatc tattcc 26

<210> 11  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 11  
gaagtatcct gggcttctga 20

<210> 12  
<211> 26  
<212> DNA  
<213> Sus scrofa

<400> 12

9/20

gtttctccag gtttccagac atccac 26

<210> 13  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 13  
gcttctgtct gccctactt 20

<210> 14  
<211> 26  
<212> DNA  
<213> Sus scrofa

<400> 14  
gtttctaagt tctactgtaa gacacc 26

<210> 15  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 15  
ccaagctgtg gtggctgaat 20

<210> 16  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 16  
cagcacagca gtgccaccta 20

<210> 17  
<211> 19  
<212> DNA  
<213> Sus scrofa

<400> 17  
caaactcttc taggcgtgt 19

<210> 18  
<211> 26  
<212> DNA  
<213> Sus scrofa

<400> 18  
gtttctggaa cttccatatg ccatgg 26

<210> 19  
<211> 20  
<212> DNA  
<213> Sus scrofa

10/20

<400> 19  
aggggtggatg gtaggcttca

20

<210> 20  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 20  
gtctcgctcc tgaaggaagt

20

<210> 21  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 21  
agtcacgtgg ccatgctatc

20

<210> 22  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 22  
ctcaactgga ttgagtcagt

20

<210> 23  
<211> 20  
<212> DNA  
<213> Sus scrofa

<400> 23  
ttggcgcaac tgttatttct

20

<210> 24  
<211> 19  
<212> DNA  
<213> Sus scrofa

<400> 24  
aggcaaagga agagcacag

19

<210> 25  
<211> 18  
<212> DNA  
<213> Sus scrofa

<400> 25  
agccgtgggc atcgttgg

18

<210> 26  
<211> 21

11/20

&lt;212&gt; DNA

&lt;213&gt; Sus scrofa

&lt;400&gt; 26

agaaggagac agacagggcga

21

&lt;210&gt; 27

&lt;211&gt; 1873

&lt;212&gt; ADN

&lt;213&gt; Sus scrofa

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1395)

&lt;400&gt; 27

atg agc ttc cta gag caa gga gag agc cgt tca tgg cca tcc cga gct	48
Met Ser Phe Leu Glu Gln Gly Glu Ser Arg Ser Trp Pro Ser Arg Ala	
1 5 10 15	
gta acc acc agc tca gaa aga agc cat ggg gac cag ggg aac aag gcc	96
Val Thr Thr Ser Ser Glu Arg Ser His Gly Asp Gln Gly Asn Lys Ala	
20 25 30	
tct aga tgg aca agg cag gag gat gta gag gaa ggg ggg cct ccg ggc	144
Ser Arg Trp Thr Arg Gln Glu Asp Val Glu Glu Gly Gly Pro Pro Gly	
35 40 45	
ccg agg gaa ggt ccc cag tcc agg cca gtt gct gag tcc acc ggg cag	192
Pro Arg Glu Gly Pro Gln Ser Arg Pro Val Ala Glu Ser Thr Gly Gln	
50 55 60	
gag gcc aca ttc ccc aag gcc aca ccc ttg gcc caa gcc gct ccc ttg	240
Glu Ala Thr Phe Pro Lys Ala Thr Pro Leu Ala Gln Ala Ala Pro Leu	
65 70 75 80	
gcc gag gtg gac aac ccc cca aca gag cgg gac atc ctc ccc tct gac	288
Ala Glu Val Asp Asn Pro Pro Thr Glu Arg Asp Ile Leu Pro Ser Asp	
85 90 95	
tgt gca gcc tca gcc tcc gac tcc aac aca gac cat ctg gat ctg ggc	336
Cys Ala Ala Ser Ala Ser Asp Ser Asn Thr Asp His Leu Asp Leu Gly	
100 105 110	
ata gag ttc tca gcc tcg gcg gcg tcg ggg gat gag ctt ggg ctg gtg	384
Ile Glu Phe Ser Ala Ser Ala Ala Ser Gly Asp Glu Leu Gly Leu Val	
115 120 125	
gaa gag aag cca gcc ccg tgc cca tcc cca gag gtg ctg tta ccc agg	432
Glu Glu Lys Pro Ala Pro Cys Pro Ser Pro Glu Val Leu Leu Pro Arg	
130 135 140	
ctg ggc tgg gat gat gag ctg cag aag ccg ggg gcc cag gtc tac atg	480
Leu Gly Trp Asp Asp Glu Leu Gln Lys Pro Gly Ala Gln Val Tyr Met	
145 150 155 160	
cac ttc atg cag gag cac acc tgc tac gat gcc atg gcg acc agc tcc	528
His Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser	
165 170 175	

12/20

aaa ctg gtc atc ttc gac acc atg ctg gag atc aag aag gcc ttc ttt	576
Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe	
180 185 190	
gcc ctg gtg gcc aac ggc gtc cga gcg gca cct ttg tgg gac agc aag	624
Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser Lys	
195 200 205	
aag cag agc ttc gtg ggg atg ctg acc atc aca gac ttc atc ttg gtg	672
Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu Val	
210 215 220	
ctg cac cgc tat tac agg tcc ccc ctg gtc cag atc tac gag att gaa	720
Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile Glu	
225 230 235 240	
gaa cat aag att gag acc tgg agg gag atc tac ctt caa ggc tgc ttc	768
Glu His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe	
245 250 255	
aag cct ctg gtc tcc atc tct ccc aat gac agc ctg ttc gaa gct gtc	816
Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val	
260 265 270	
tac gcc ctc atc aag aac cgg atc cac cgc ctg ccg gtc ctg gac cct	864
Tyr Ala Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp Pro	
275 280 285	
gtc tcc ggg gct gtg ctc cac atc ctc aca cat aag cgg ctt ctc aag	912
Val Ser Gly Ala Val Leu His Ile Leu Thr His Lys Arg Leu Leu Lys	
290 295 300	
ttc ctg cac atc ttt ggc acc ctg ctg ccc cgg ccc tcc ttc ctc tac	960
Phe Leu His Ile Phe Gly Thr Leu Leu Pro Arg Pro Ser Phe Leu Tyr	
305 310 315 320	
cgc acc atc caa gat ttg ggc atc ggc aca ttc cga gac ttg gcc gtg	1008
Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala Val	
325 330 335	
gtg ctg gaa acg gcg ccc atc ctg acc gca ctg gac atc ttc gtg gac	1056
Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val Asp	
340 345 350	
cgg cgt gtg tct gcg ctg cct gtg gtc aac gaa act gga cag gta gtg	1104
Arg Arg Val Ser Ala Leu Pro Val Val Asn Glu Thr Gly Gln Val Val	
355 360 365	
ggc ctc tac tct cgc ttt gat gtg atc cac ctg gct gcc caa caa aca	1152
Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln Thr	
370 375 380	
tac aac cac ctg gac atg aat gtg gga gaa gcc ctg agg cag cgg aca	1200
Tyr Asn His Leu Asp Met Asn Val Gly Glu Ala Leu Arg Gln Arg Thr	
385 390 395 400	
ctg tgt ctg gaa ggc gtc ctt tcc tgc cag ccc cac gag acc ttg ggg	1248
Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Thr Leu Gly	
405 410 415	

13/20

gaa gtc att gac cgg att gtc cgg gaa cag gtg cac cgc ctg gtg ctc 1296  
 Glu Val Ile Asp Arg Ile Val Arg Glu Gln Val His Arg Leu Val Leu  
                   420                                  425                                  430

gtg gat gag acc cag cac ctt ctg ggc gtg gtg tcc ctc tct gac atc 1344  
 Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp Ile  
                   435                                  440                                  445

ctt cag gct ctg gtg ctc agc cct gct gga att gat gcc ctc ggg gcc 1392  
 Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly Ala  
                   450                                  455                                  460

tga gaaccttgga acctttgctc tcaggccacc tggcacacct ggaagccagt 1445  
 465

gaagggagcc gtggactcag ctctcacttc ccctcagccc cacttgctgg tctggctctt 1505  
 gttcaggtag gctccgcccc gggccccctgg cctcagcacc agccccctcag tctccctggg 1565  
 caccagatc tcagactggg gcacctgaa gatgggagtg gccagctta tagctgagca 1625  
 gccttgtagaa atctaccagc atcaagactc actgtgggac cactgctttg tcccattctc 1685  
 agctgaaatg atggagggcc tcataagagg ggtggacagg gcctggagta gaggccagat 1745  
 cagtgacgtg ccttcaggac ctccggggag ttagagctgc cctctctcag ttcagttccc 1805  
 ccctgctgag aatgtccctg gaaggaagcc agttaataaa ccttggttgg atggaatttg 1865  
 gagagtcg 1873

<210> 28  
 <211> 464  
 <212> PRT  
 <213> Sus scrofa

<400> 28  
 Met Ser Phe Leu Glu Gln Gly Glu Ser Arg Ser Trp Pro Ser Arg Ala  
   1                                  5                                  10                                  15  
 Val Thr Thr Ser Ser Glu Arg Ser His Gly Asp Gln Gly Asn Lys Ala  
                                   20                                  25                                  30  
 Ser Arg Trp Thr Arg Gln Glu Asp Val Glu Glu Gly Gly Pro Pro Gly  
                                   35                                  40                                  45  
 Pro Arg Glu Gly Pro Gln Ser Arg Pro Val Ala Glu Ser Thr Gly Gln  
                                   50                                  55                                  60  
 Glu Ala Thr Phe Pro Lys Ala Thr Pro Leu Ala Gln Ala Ala Pro Leu  
                                   65                                  70                                  75                                  80  
 Ala Glu Val Asp Asn Pro Pro Thr Glu Arg Asp Ile Leu Pro Ser Asp  
                                   85                                  90                                  95  
 Cys Ala Ala Ser Ala Ser Asp Ser Asn Thr Asp His Leu Asp Leu Gly  
                                   100                                  105                                  110  
 Ile Glu Phe Ser Ala Ser Ala Ala Ser Gly Asp Glu Leu Gly Leu Val  
                                   115                                  120                                  125  
 Glu Glu Lys Pro Ala Pro Cys Pro Ser Pro Glu Val Leu Leu Pro Arg  
                                   130                                  135                                  140  
 Leu Gly Trp Asp Asp Glu Leu Gln Lys Pro Gly Ala Gln Val Tyr Met  
  145                                  150                                  155                                  160  
 His Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser  
                                   165                                  170                                  175

SUBSTITUTE SHEET (RULE 26)



14/20

Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe  
 180 185 190  
 Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser Lys  
 195 200 205  
 Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu Val  
 210 215 220  
 Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile Glu  
 225 230 235 240  
 Glu His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe  
 245 250 255  
 Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val  
 260 265 270  
 Tyr Ala Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp Pro  
 275 280 285  
 Val Ser Gly Ala Val Leu His Ile Leu Thr His Lys Arg Leu Leu Lys  
 290 295 300  
 Phe Leu His Ile Phe Gly Thr Leu Leu Pro Arg Pro Ser Phe Leu Tyr  
 305 310 315 320  
 Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala Val  
 325 330 335  
 Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val Asp  
 340 345 350  
 Arg Arg Val Ser Ala Leu Pro Val Val Asn Glu Thr Gly Gln Val Val  
 355 360 365  
 Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln Thr  
 370 375 380  
 Tyr Asn His Leu Asp Met Asn Val Gly Glu Ala Leu Arg Gln Arg Thr  
 385 390 395 400  
 Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Thr Leu Gly  
 405 410 415  
 Glu Val Ile Asp Arg Ile Val Arg Glu Gln Val His Arg Leu Val Leu  
 420 425 430  
 Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp Ile  
 435 440 445  
 Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly Ala  
 450 455 460

<210> 29  
 <211> 2115  
 <212> ADN  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(1395)

<400> 29  
 atg agc ttc cta gag caa gaa aac agc agc tca tgg cca tca cca gct 48  
 Met Ser Phe Leu Glu Gln Glu Asn Ser Ser Ser Trp Pro Ser Pro Ala  
 1 5 10 15  
 gtg acc agc agc tca gaa aga atc cgt ggg aaa cgg agg gcc aaa gcc 96  
 Val Thr Ser Ser Ser Glu Arg Ile Arg Gly Lys Arg Arg Ala Lys Ala  
 20 25 30  
 ttg aga tgg aca agg cag aag tcg gtg gag gaa ggg gag cca cca ggt 144  
 Leu Arg Trp Thr Arg Gln Lys Ser Val Glu Glu Gly Glu Pro Pro Gly  
 35 40 45

15/20

cag ggg gaa ggt ccc cgg tcc agg cca act gct gag tcc acc ggg ctg	192
Gln Gly Glu Gly Pro Arg Ser Arg Pro Thr Ala Glu Ser Thr Gly Leu	
50 55 60	
gag gcc aca ttc ccc aag acc aca ccc ttg gct caa gct gat cct gcc	240
Glu Ala Thr Phe Pro Lys Thr Thr Pro Leu Ala Gln Ala Asp Pro Ala	
65 70 75 80	
ggg gtg ggc act cca cca aca ggg tgg gac tgc ctc ccc tct gac tgt	288
Gly Val Gly Thr Pro Pro Thr Gly Trp Asp Cys Leu Pro Ser Asp Cys	
85 90 95	
aca gcc tca gct gca ggc tcc agc aca gat gat gtg gag ctg gcc acg	336
Thr Ala Ser Ala Ala Gly Ser Ser Thr Asp Asp Val Glu Leu Ala Thr	
100 105 110	
gag ttc cca gcc aca gag gcc tgg gag tgt gag cta gaa ggc ctg ctg	384
Glu Phe Pro Ala Thr Glu Ala Trp Glu Cys Glu Leu Glu Gly Leu Leu	
115 120 125	
gaa gag agg cct gcc ctg tgc ctg tcc ccg cag gcc cca ttt ccc aag	432
Glu Glu Arg Pro Ala Leu Cys Leu Ser Pro Gln Ala Pro Phe Pro Lys	
130 135 140	
ctg ggc tgg gat gac gaa ctg cgg aaa ccc ggc gcc cag atc tac atg	480
Leu Gly Trp Asp Asp Glu Leu Arg Lys Pro Gly Ala Gln Ile Tyr Met	
145 150 155 160	
cgc ttc atg cag gag cac acc tgc tac gat gcc atg gca act agc tcc	528
Arg Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser	
165 170 175	
aag cta gtc atc ttc gac acc atg ctg gag atc aag aag gcc ttc ttt	576
Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe	
180 185 190	
gct ctg gtg gcc aac ggt gtg cgg gca gcc cct cta tgg gac agc aag	624
Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser Lys	
195 200 205	
aag cag agc ttt gtg ggg atg ctg acc atc act gac ttc atc ctg gtg	672
Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu Val	
210 215 220	
ctg cat cgc tac tac agg tcc ccc ctg gtc cag atc tat gag att gaa	720
Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile Glu	
225 230 235 240	
caa cat aag att gag acc tgg agg gag atc tac ctg caa ggc tgc ttc	768
Gln His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe	
245 250 255	
aag cct ctg gtc tcc atc tct cct aat gat agc ctg ttt gaa gct gtc	816
Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val	
260 265 270	
tac acc ctc atc aag aac cgg atc cat cgc ctg cct gtt ctt gac ccg	864
Tyr Thr Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp Pro	
275 280 285	

16/20

gtg tca ggc aac gta ctc cac atc ctc aca cac aaa cgc ctg ctc aag 912  
 Val Ser Gly Asn Val Leu His Ile Leu Thr His Lys Arg Leu Leu Lys  
 290 295 300

ttc ctg cac atc ttt ggt tcc ctg ctg ccc cgg ccc tcc ttc ctc tac 960  
 Phe Leu His Ile Phe Gly Ser Leu Leu Pro Arg Pro Ser Phe Leu Tyr  
 305 310 315 320

cgc act atc caa gat ttg ggc atc ggc aca ttc cga gac ttg gct gtg 1008  
 Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala Val  
 325 330 335

gtg ctg gag aca gca ccc atc ctg act gca ctg gac atc ttt gtg gac 1056  
 Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val Asp  
 340 345 350

cgg cgt gtg tct gca ctg cct gtg gtc aac gaa tgt ggt cag gtc gtg 1104  
 Arg Arg Val Ser Ala Leu Pro Val Val Asn Glu Cys Gly Gln Val Val  
 355 360 365

ggc ctc tat tcc cgc ttt gat gtg att cac ctg gct gcc cag caa acc 1152  
 Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln Thr  
 370 375 380

tac aac cac ctg gac atg agt gtg gga gaa gcc ctg agg cag agg aca 1200  
 Tyr Asn His Leu Asp Met Ser Val Gly Glu Ala Leu Arg Gln Arg Thr  
 385 390 395 400

cta tgt ctg gag gga gtc ctt tcc tgc cag ccc cac gag agc ttg ggg 1248  
 Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Ser Leu Gly  
 405 410 415

gaa gtg atc gac agg att gct cgg gag cag gta cac agg ctg gtg cta 1296  
 Glu Val Ile Asp Arg Ile Ala Arg Glu Gln Val His Arg Leu Val Leu  
 420 425 430

gtg gac gag acc cag cat ctc ttg ggc gtg gtc tcc ctc tcc gac atc 1344  
 Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp Ile  
 435 440 445

ctt cag gca ctg gtg ctc agc cct gct ggc atc gat gcc ctc ggg gcc 1392  
 Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly Ala  
 450 455 460

tga gaagatctga gtcctcaatc ccaagccaac tgcacactgg aagccaatga 1445  
 465

aggaattgag aacagcttca tttccccaac cccaatttgc tgggttcagct atgattcagg 1505

cttcttcagc cttccaaaat tgcctttgcc ttacttgtgc tcccagaacc cttcgggcat 1565

gcccagtga ccatgggatg atgaaattaa ggagaacagc tgagtcaagc ttggagggtcc 1625

ctgaaccaga ggcactagga ttaccccagg gccatctgtg ctccatgcc gcccatcccc 1685

ttgccgcctg actggggtcgg atggccccag tgggtttagt cagggcttct ggattcctcg 1745

gtttctgggc tacctatggc ttacgccttc agctcctggg agtcccagct gttgttccca 1805

gcaacgtcgc cactgccctc ctactctcca ggctttgtca tttcaaggct gctgaaatgc 1865

17/20

tgcatttcag gggccaccat ggagcagccg ttatttatag aactgcctgt tggaggtggg 1925  
 gagtcctccc tccattcttg tccagaaaac tccttagctc tcgcagtgag ccatgttctt 1985  
 agtctccagg gatggatggc cttgtatatg gacccttgag aatgagcaat tgagaaaaca 2045  
 aaacaaaagg aacaatccat gaacttagat tttattgggt tcaactcaaaa tgctgcagtc 2105  
 atttgacctg 2115

<210> 30  
 <211> 464  
 <212> PRT  
 <213> Homo sapiens

<400> 30  
 Met Ser Phe Leu Glu Gln Glu Asn Ser Ser Ser Trp Pro Ser Pro Ala  
 1 5 10 15  
 Val Thr Ser Ser Ser Glu Arg Ile Arg Gly Lys Arg Arg Ala Lys Ala  
 20 25 30  
 Leu Arg Trp Thr Arg Gln Lys Ser Val Glu Glu Gly Glu Pro Pro Gly  
 35 40 45  
 Gln Gly Glu Gly Pro Arg Ser Arg Pro Thr Ala Glu Ser Thr Gly Leu  
 50 55 60  
 Glu Ala Thr Phe Pro Lys Thr Thr Pro Leu Ala Gln Ala Asp Pro Ala  
 65 70 75 80  
 Gly Val Gly Thr Pro Pro Thr Gly Trp Asp Cys Leu Pro Ser Asp Cys  
 85 90 95  
 Thr Ala Ser Ala Ala Gly Ser Ser Thr Asp Asp Val Glu Leu Ala Thr  
 100 105 110  
 Glu Phe Pro Ala Thr Glu Ala Trp Glu Cys Glu Leu Glu Gly Leu Leu  
 115 120 125  
 Glu Glu Arg Pro Ala Leu Cys Leu Ser Pro Gln Ala Pro Phe Pro Lys  
 130 135 140  
 Leu Gly Trp Asp Asp Glu Leu Arg Lys Pro Gly Ala Gln Ile Tyr Met  
 145 150 155 160  
 Arg Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr Ser Ser  
 165 170 175  
 Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala Phe Phe  
 180 185 190  
 Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp Ser Lys  
 195 200 205  
 Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile Leu Val  
 210 215 220  
 Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu Ile Glu  
 225 230 235 240  
 Gln His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly Cys Phe  
 245 250 255  
 Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu Ala Val  
 260 265 270  
 Tyr Thr Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu Asp Pro  
 275 280 285  
 Val Ser Gly Asn Val Leu His Ile Leu Thr His Lys Arg Leu Leu Lys  
 290 295 300  
 Phe Leu His Ile Phe Gly Ser Leu Leu Pro Arg Pro Ser Phe Leu Tyr  
 305 310 315 320  
 Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu Ala Val  
 325 330 335

18/20

Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe Val Asp  
 340 345 350  
 Arg Arg Val Ser Ala Leu Pro Val Val Asn Glu Cys Gly Gln Val Val  
 355 360 365  
 Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln Gln Thr  
 370 375 380  
 Tyr Asn His Leu Asp Met Ser Val Gly Glu Ala Leu Arg Gln Arg Thr  
 385 390 395 400  
 Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Ser Leu Gly  
 405 410 415  
 Glu Val Ile Asp Arg Ile Ala Arg Glu Gln Val His Arg Leu Val Leu  
 420 425 430  
 Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser Asp Ile  
 435 440 445  
 Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu Gly Ala  
 450 455 460

&lt;210&gt; 31

&lt;211&gt; 2022

&lt;212&gt; ADN

&lt;213&gt; Sus scrofa

&lt;400&gt; 31

atggagcttg ccgagctaga gcaggcactg cgcagggtcc cgggggtccc ggggggctgg 60  
 gagctggagc aactgaggcc agagggcaga gggcccacca ctgctggatac tccctcctgg 120  
 agcagcctcg ggggaccta gcatcaagag atgagcttcc tagagcaagg agagagccgt 180  
 tcatggccat cccgagctgt aaccaccagc tcagaaagaa gccatgggga ccaggggaac 240  
 aaggcctcta gatggacaag gcaggaggat gtagaggaag gggggcctcc gggcccgagg 300  
 gaaggtcccc agtccaggcc agttgctgag tccaccgggc aggaggccac attccccaa 360  
 gccacacctt tggcccaagc cgctcccttg gccgaggtgg acaaccccc aacagagcgg 420  
 gacatcctcc cctctgactg tgcagcctca gcctccgact ccaacacaga ccctctggat 480  
 ctgggcatag agttctcagc ctggcgggcg tcgggggatg agcttgggct ggtggaagag 540  
 aagccagccc cgtgccatc cccagaggtg ctgttaccca ggctgggctg ggatgatgag 600  
 ctgcagaagc cgggggcccc ggtctacatg cacttcacgc aggagcacac ctgctacgat 660  
 gccatggcga ccagctcaa actggtcatc ttcgacacca tgctggagat caagaaggcc 720  
 ttctttgccc tgggtggcaa cggcgtccga gcggcacctt tgtgggacag caagaagcag 780  
 agcttcgtgg ggatgctgac catcacagac ttcattcttg tgctgcaccg ctattacagg 840  
 tccccctgg tccagatcta cgagattgaa gaacataaga ttgagacctg gagggagatc 900  
 taccttcaag gctgcttcaa gcctctggtc tccatctctc ccaatgacag cctgttcgaa 960  
 gctgtctacg ccctcatcaa gaaccggatc caccgcctgc cggctcctgga ccctgtctcc 1020  
 ggggctgtgc tccacatcct cacacataag cggcttctca agttcctgca catctttggc 1080  
 accctgctgc cccggccctc ctctctctac cgcaccatcc aagatttggg catcggcaca 1140  
 ttccgagact tggccgtggg gctggaaacg gcgcccaccc tgaccgcact ggacatcttc 1200  
 gtggaccggc gtgtgtctgc gctgctgtg gtcaacgaaa ctggacaggt agtgggcctc 1260  
 tactctcgtc ttgatgtgat ccacctggct gcccaacaaa catacaacca cctggacatg 1320  
 aatgtgggag aagccctgag gcagcggaca ctgtgtctgg aaggcgtcct ttctgcccag 1380  
 cccacagaga ccttggggga agtcattgac cggattgtcc gggaacaggt gcaccgcctg 1440  
 gtgctcgtgg atgagaccca gcacctctg ggcgtggtgt ccctctctga catccttcag 1500  
 gctctggtgc tcagccctgc tggaaattgat gccctcgggg cctgagaacc ttggaacctt 1560  
 tgctctcagg ccacctggca cacctggaag ccagtgaagg gagccgtgga ctcagctctc 1620  
 acttccccct agccccactt gctggctctg ctctgttca ggtaggctcc gccgggggcc 1680  
 cctggcctca gcatcagccc ctgagctctc ctgggcaccc agatctcaga ctggggcacc 1740  
 ctgaagatgg gagtggccca gcttatagct gagcagcctt gtgaaatcta ccagcatcaa 1800  
 gactcactgt gggaccactg ctttgtccca ttctcagctg aaatgatgga gggcctcata 1860  
 agaggggtgg acagggcctg gtagtagggc cagatcagtg acgtgccttc agggacctccg 1920  
 gggagttaga gctgccctct ctcagttcag ttccccctg ctgagaatgt ccctggaagg 1980  
 aagccagtta ataaacctt gttggatgga atttggagag tc 2022

19/20

<210> 32  
 <211> 514  
 <212> PRT  
 <213> Sus scrofa

<400> 32  
 Met Glu Leu Ala Glu Leu Glu Gln Ala Leu Arg Arg Val Pro Gly Ser  
   1                  5                  10                  15  
 Arg Gly Gly Trp Glu Leu Glu Gln Leu Arg Pro Glu Gly Arg Gly Pro  
           20                  25                  30  
 Thr Thr Ala Asp Thr Pro Ser Trp Ser Ser Leu Gly Gly Pro Lys His  
           35                  40                  45  
 Gln Glu Met Ser Phe Leu Glu Gln Gly Glu Ser Arg Ser Trp Pro Ser  
   50                  55                  60  
 Arg Ala Val Thr Thr Ser Ser Glu Arg Ser His Gly Asp Gln Gly Asn  
 .65                  70                  75                  80  
 Lys Ala Ser Arg Trp Thr Arg Gln Glu Asp Val Glu Glu Gly Gly Pro  
           85                  90                  95  
 Pro Gly Pro Arg Glu Gly Pro Gln Ser Arg Pro Val Ala Glu Ser Thr  
           100                  105                  110  
 Gly Gln Glu Ala Thr Phe Pro Lys Ala Thr Pro Leu Ala Gln Ala Ala  
   115                  120                  125  
 Pro Leu Ala Glu Val Asp Asn Pro Pro Thr Glu Arg Asp Ile Leu Pro  
   130                  135                  140  
 Ser Asp Cys Ala Ala Ser Ala Ser Asp Ser Asn Thr Asp His Leu Asp  
 145                  150                  155                  160  
 Leu Gly Ile Glu Phe Ser Ala Ser Ala Ala Ser Gly Asp Glu Leu Gly  
           165                  170                  175  
 Leu Val Glu Glu Lys Pro Ala Pro Cys Pro Ser Pro Glu Val Leu Leu  
           180                  185                  190  
 Pro Arg Leu Gly Trp Asp Asp Glu Leu Gln Lys Pro Gly Ala Gln Val  
           195                  200                  205  
 Tyr Met His Phe Met Gln Glu His Thr Cys Tyr Asp Ala Met Ala Thr  
   210                  215                  220  
 Ser Ser Lys Leu Val Ile Phe Asp Thr Met Leu Glu Ile Lys Lys Ala  
 225                  230                  235                  240  
 Phe Phe Ala Leu Val Ala Asn Gly Val Arg Ala Ala Pro Leu Trp Asp  
           245                  250                  255  
 Ser Lys Lys Gln Ser Phe Val Gly Met Leu Thr Ile Thr Asp Phe Ile  
           260                  265                  270  
 Leu Val Leu His Arg Tyr Tyr Arg Ser Pro Leu Val Gln Ile Tyr Glu  
   275                  280                  285

SUBSTITUTE SHEET (RULE 26)

20/20

Ile Glu Glu His Lys Ile Glu Thr Trp Arg Glu Ile Tyr Leu Gln Gly  
 290 295 300  
 Cys Phe Lys Pro Leu Val Ser Ile Ser Pro Asn Asp Ser Leu Phe Glu  
 305 310 315 320  
 Ala Val Tyr Ala Leu Ile Lys Asn Arg Ile His Arg Leu Pro Val Leu  
 325 330 335  
 Asp Pro Val Ser Gly Ala Val Leu His Ile Leu Thr His Lys Arg Leu  
 340 345 350  
 Leu Lys Phe Leu His Ile Phe Gly Thr Leu Leu Pro Arg Pro Ser Phe  
 355 360 365  
 Leu Tyr Arg Thr Ile Gln Asp Leu Gly Ile Gly Thr Phe Arg Asp Leu  
 370 375 380  
 Ala Val Val Leu Glu Thr Ala Pro Ile Leu Thr Ala Leu Asp Ile Phe  
 385 390 395 400  
 Val Asp Arg Arg Val Ser Ala Leu Pro Val Val Asn Gly Thr Gly Gln  
 405 410 415  
 Val Val Gly Leu Tyr Ser Arg Phe Asp Val Ile His Leu Ala Ala Gln  
 420 425 430  
 Gln Thr Tyr Asn His Leu Asp Met Asn Val Gly Glu Ala Leu Arg Gln  
 435 440 445  
 Arg Thr Leu Cys Leu Glu Gly Val Leu Ser Cys Gln Pro His Glu Thr  
 450 455 460  
 Leu Gly Glu Val Ile Asp Arg Ile Val Arg Glu Gln Val His Arg Leu  
 465 470 475 480  
 Val Leu Val Asp Glu Thr Gln His Leu Leu Gly Val Val Ser Leu Ser  
 485 490 495  
 Asp Ile Leu Gln Ala Leu Val Leu Ser Pro Ala Gly Ile Asp Ala Leu  
 500 505 510  
 Gly Ala

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number  
**WO 01/20003 A3**

(51) International Patent Classification?: C12N 15/54,  
15/11, 9/12, C12Q 1/68, A01K 67/027, G01N 33/68

(21) International Application Number: PCT/EP00/09896

(22) International Filing Date:  
11 September 2000 (11.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
99402236.6 10 September 1999 (10.09.1999) EP  
00401388.4 18 May 2000 (18.05.2000) EP

(71) Applicant (for all designated States except US):  
**INSTITUT NATIONAL DE LA RECHERCHE  
AGRONOMIQUE (INRA)** [FR/FR]; 147, rue de l'Uni-  
versité, F-75007 Paris (FR).

(71) Applicants and

(72) Inventors: **ANDERSSON, Leif** [SE/SE]; Bergagatan 30,  
S-752 39 Uppsala (SE). **LOOFT, Christian** [DE/DE];  
Mittelweg 3 c, 24802 Bokelholm (DE). **KALM, Ernst**  
[DE/DE]; Schmalholt 1, 24239 Achterwehr (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MILAN, Denis**  
[FR/FR]; 3 bis, chemin du Tricou, F-31670 Labège (FR).

**RORIC, Annie** [FR/FR]; 33, rue des Capitouls, F-31650  
Saint-Orens-De-Gameville (FR). **ROGEL-GAILLARD,**  
**Claire** [FR/FR]; 156, rue Léon Maurice Nordmann,  
F-75013 Paris (FR). **IANNUCELLI, Nathalie** [FR/FR];  
7, boulevard des Alouettes, F-31320 Castanet-Tolosan  
(FR). **GELLIN, Joël** [FR/FR]; 8, allée des Amazones,  
F-31320 Auzeville (FR). **LE ROY, Pascale** [FR/FR]; 32,  
avenue Saint Marc, F-91300 Massy (FR). **CHARDON,**  
**Patrick** [FR/FR]; 17, rue de Petite Fontaine, F-91430  
Vauhallan (FR).

(74) Agents: **VIALLE-PRESLES, Marie-José et al.**; Cabinet  
Ores, 6, avenue de Messine, F-75008 Paris (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

[Continued on next page]

(54) Title: VARIANTS OF THE GAMMA CHAIN OF AMPK, DNA SEQUENCES ENCODING THE SAME, AND USES THEREOF



(57) Abstract: The invention concerns variants of the gamma chain of vertebrate AMP-activated kinase (AMPK), as well as nucleic acid sequences encoding said variants and use thereof for the diagnosis or treatment of dysfunction of energy metabolisms.





— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(88) Date of publication of the international search report:  
17 May 2001

## INTERNATIONAL SEARCH REPORT

International Application No.

PC./EP 00/09896

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/11 C12N9/12 C12Q1/68 A01K67/027  
G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q A01K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HILLIER ET AL.: "WashU-NCI human EST project" EMBL DATABASE ACC NO: AA178898, 1 January 1997 (1997-01-01), XP002130593 cited in the application abstract	1-5, 11-17
X	ROBIC ET AL.: "A radiation hybrid map of the RN region in pigs demonstrates conserved gene order compared with the human and mouse genomes" MAMMALIAN GENOME, vol. 10, no. 6, June 1999 (1999-06), pages 565-568, XP000876695 cited in the application page 565	29-33
A	page 567; figure 1; table 1 --- -/-	1-28, 32-37

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

7 March 2001

Date of mailing of the international search report

19/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
Fax: (+31-70) 340-3016

Authorized officer

van Klompenburg, W

## INTERNATIONAL SEARCH REPORT

International Application No

PC../EP 00/09896

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category -	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 25341 A (ST VINCENTS INST MED RES ;DARTMOUTH COLLEGE (US); KEMP BRUCE E (AU) 17 July 1997 (1997-07-17) page 18, line 21 - line 34 page 25, line 1 -page 28, line 5 claims 16-21	13-15
A	---	1-12, 16-37
X	WO 98 58052 A (INCYTE PHARMA INC ;CORLEY NEIL C (US); BANDMAN OLGA (US); GOLI SUR) 23 December 1998 (1998-12-23) SEQ ID NOs 7 & 14 page 18, line 21 -page 19, line 8 page 38, line 6 -page 44, line 29 claims 1-24; figure 7	13-15
A	---	1-12, 16-37
X	WATERSTON: "Homo sapiens chromosome unknown clone NH0459I19" EMBL DATABASE ACC NO: AC009974, 9 September 1999 (1999-09-09), XP002130594 abstract	11-16
X	---	
X	HILLIER ET AL.: "The WashU-Merck EST Project" EMBL DATABASE ACC NO: W94830, 17 July 1996 (1996-07-17), XP002130595 abstract	11-16
A	---	
A	MILAN ET AL.: "Accurate mapping of the "acid meat" RN gene on genetic and physical maps of pig chromosome 15" MAMMALIAN GENOME, vol. 7, no. 1, January 1996 (1996-01), pages 47-51, XP000876743 cited in the application page 50, column 2 -page 51, column 1; figure 1	1-35
	---	
	-/--	

## INTERNATIONAL SEARCH REPORT

International Application No

PC./EP 00/09896

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>CHEUNG ET AL.: "Characterization of AMP-activated protein kinase gamma-subunit isoforms and their role in AMP binding" BIOCHEMICAL JOURNAL, vol. 346, no. 3, 15 March 2000 (2000-03-15), pages 659-669, XP002162237 figures 2,3,5; tables 1,2 -&amp; DATABASE EMBL 'Online!' EBI; ACC. NO.: AJ249977, 7 January 2000 (2000-01-07)</p> <p>CARLING : "Homo sapiens mRNA for AMP-activated protein kinase gamma 3 subunit (AMPK gamma 3 gene)" XP002162239 abstract</p>	1-4, 11-17, 20,37
P,X	<p>MILAN ET AL.: "A mutation in PRKAG3 associated with excess glycogen content in pig skeletal muscle" SCIENCE, vol. 288, 19 May 2000 (2000-05-19), pages 1248-1251, XP002162238 figures 1,2; tables 1-3</p>	1-33

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/09896

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9725341	A	17-07-1997	AU 714905 B	13-01-2000
			AU 1693697 A	01-08-1997
			CA 2241786 A	17-07-1997
			EP 0873354 A	28-10-1998
			JP 2000503202 T	21-03-2000
			US 6124125 A	26-09-2000
WO 9858052	A	23-12-1998	US 5885803 A	23-03-1999
			AU 8154798 A	04-01-1999
			EP 1007692 A	14-06-2000